

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

Does adrenal responsiveness vary with sex and reproductive status in  
*Egernia whitii*, a viviparous skink?

Victoria A. Cartledge\* and Susan M. Jones

School of Zoology, University of Tasmania, Private Bag 5, Hobart TAS 7001,  
Australia

<sup>a</sup> current address: Zoology, School of Animal Biology, The University of Western  
Australia, 35 Stirling Highway, Crawley, WA 6009, Australia

\* Corresponding author. Tel.: + 61-8-6488-1483; fax: + 61-8-6488-1029  
*E-mail address:* vcartled@cyllene.uwa.edu.au (V. A. Cartledge)

1 **Abstract**

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

In mammals, oestrogens generally stimulate adrenal function whilst androgens are inhibitory, and gestating females down-regulate their acute response to stressors in order to protect current reproductive investment. This study aimed to determine if adrenocortical function is similarly modulated by sex and reproductive status in the viviparous lizard, *Egernia whitii*. We compared the adrenocortical response to acute capture stress in female *E. whitii* during active (post-ovulatory and gestating) and quiescent (post-partum) phases of their reproductive cycle. We also compared the responses of reproductively quiescent males and females to acute stress and ACTH challenge to determine if there are sex-related differences in HPA axis activity when the influence of reproductive hormones is minimal. The females' responses to acute capture stress varied significantly with reproductive stage, and quiescent females displayed the strongest immediate response, with a rapid and sustained increase in plasma corticosterone (CORT) concentrations. Post-ovulatory females showed the most conservative adrenocortical response and while gestating females showed a large immediate response, this was not as prolonged as in quiescent females. Reproductively quiescent males and females exhibited similar responses to acute stress, and also responded similarly to ACTH injection, with plasma CORT reaching maximal concentrations of 52.1 ng/mL and 59.4 ng/mL respectively. Reproductively quiescent females treated with oestrogen exhibited greater responsiveness to ACTH than males, although basal plasma CORT concentrations were unaltered: these results suggest that the attenuation of the acute stress response observed in reproductively active females of *E. whitii* may be regulated upstream of ACTH secretion. Our results demonstrate that the activity of the HPA axis is modulated by reproductive status in this viviparous reptile, and that gestating females are able to buffer their embryos from the potentially adverse effects of elevated plasma corticosteroids.

*Keywords: ACTH, Capture, Corticosterone, Oestrogen, Reptile, Stress, Viviparity*

## 1. Introduction

In vertebrates, the hypothalamo-pituitary-adrenal (HPA) axis is central to promoting survival by allowing the animal to respond behaviourally and physiologically to environmental stressors. Such responses may, however, include suppression of reproductive activity via inhibition of the hypothalamo-pituitary-gonadal (HPG) axis: current survival is then traded off against future reproductive success (Klose et al., 2006; Wingfield, 2003). Modulation of adrenocortical activity during particular stages of life history may therefore be an important factor in determining an animal's overall fitness (Moore and Jessop, 2003). For example, in arctic (Astheimer et al., 1995; Wingfield et al., 1995) and in desert (Wingfield et al., 1992) birds with limited time for breeding, the stress response is suppressed during the reproductive phase and breeding appears to be prioritised, albeit at the risk of increased mortality due to decreased responses to immediate stressors.

The anti-gonadal effects of stress in reptiles have been well documented (e.g. Lance and Elsey, 1986; Greenberg and Winfield, 1987; Mahmoud et al., 1989; Elsey et al., 1991, Moore et al., 1991), but reciprocal influences of reproductive status on the function of the HPA axis have received much less attention. Marine turtles have been the most comprehensively studied group, with the consistent finding that females exhibit a depressed adrenocortical response to stress when breeding (*Lepidochelys mydas*: Valverde et al., 1999; *Chelonia mydas* and *Eretmochelys imbricata*: Jessop, 2001). This appears to represent a mechanism to ensure that the large energetic reserves accumulated by these capital (oviparous) breeders are not wasted through CORT-mediated stress responses (Jessop, 2001). Similarly in female tree lizards, *Urosaurus ornatus*, vitellogenic females show a large increase in plasma CORT concentrations in response to an acute stressor, but gravid females do not (Woodley and Moore, 2002). In the garter snake *Thamnophis sirtalis*, breeding males from populations with shorter breeding seasons exhibit reduced adrenocortical responses to stress compared with males from latitudes allowing more extended breeding (Moore et al., 2001), reflecting the greater potential of stressors to affect reproductive success in males with shorter breeding seasons.

In mammals, both basal and stress induced concentrations of glucocorticoids are greater in females than in males (Gaskin and Kitay, 1970; Handa et al., 1994a). Sex differences in stress responses of reptiles remain largely unknown but there is some evidence to suggest that reproductive steroids may influence stress responses, with females exhibiting greater adrenocortical responses to stressors than males. For example, Grassman and Hess (1992a) examined the effect of gonadectomy on the acute stress responses of male and female lizards of the species *Cnemidophorus sexlineatus*: intact females subject to the acute stress of hand-chasing in a pail exhibit higher plasma CORT concentrations than males exposed to the same stressor, while castrated males show more dramatic increases in plasma CORT concentrations following acute stress than intact males. In marine green and hawksbill turtles, differences in the stress responses of females and males of differing reproductive states are evident within the first few hours (Jessop, 2001). It remains unclear, however, whether there are differences in the activation of the HPA axis independent of gonadal hormone effects.

1 Adrenal function may also vary between reproductive stages. In mammals,  
2 there is strong evidence that gestating females down-regulate their acute stress  
3 responses to protect their current reproductive investment (Neumann et al., 1998), and  
4 the negative effects of high maternal corticosteroids on the development of the foetus  
5 have been repeatedly demonstrated (e.g. Hansen et al., 1999; Weinstock, 2001). Do  
6 viviparous female reptiles similarly buffer their developing embryos against the  
7 potentially detrimental effects of high levels of circulating corticosteroids? The  
8 available evidence seems contradictory. Pregnant females of the viviparous gecko  
9 *Hoplodactylus maculatus* exhibit no suppression of the stress response to capture  
10 compared with vitellogenic females although both groups do exhibit greater stress  
11 responses than males (Cree et al., 2003), and there is no observable effect on offspring  
12 quality in females of this species challenged with ACTH during gestation (Preest et  
13 al., 2005). Gestating and post-partum females of the viviparous skink *Egernia whitii*  
14 show very similar adrenocortical responses to chronic captivity with corticosterone  
15 concentrations returning to baseline levels 1 wk following capture (Cartledge et al.,  
16 2005). However, the acute stress response has not been compared between  
17 reproductive states in this species and it may be more adaptive to modulate acute  
18 stress responses than chronic ones. Once a stressor becomes chronic in nature the  
19 reproductive benefits of curtailing the adrenocortical response are likely to be  
20 outweighed by the potential threat to fitness or mortality due to the chronic stressor.  
21

22 Thus although there appears to be considerable inter-species variation in how,  
23 and if, stress responses are modulated with reproductive status, there is evidence of  
24 reciprocal relationships between the adrenal and gonadal axes in reptiles. Such results  
25 may reflect variations in circulating concentrations of gonadal steroids. Generally,  
26 oestrogens are stimulatory to adrenal function and androgens are inhibitory (see  
27 Handa et al., 1994a for review). For example, in female rats, increased basal CORT  
28 concentrations, as well as increased ACTH and CORT responses to stressors, have  
29 been correlated with higher oestradiol concentrations (Kitay, 1963; Viau and Meaney,  
30 1991; Burgess and Handa, 1992), but there is only indirect evidence that oestrogens  
31 may stimulate adrenal activity in reptiles. In female *Uta stansburiana* lizards, peak  
32 CORT concentrations are correlated with increased gonadal mass (Wilson and  
33 Wingfield, 1992), and in females of the viviparous *Lacerta vivipara*, plasma CORT  
34 varies significantly with reproductive status, while no significant variation was  
35 observed for males (Dauphin-Villemant et al., 1990).  
36

37 Building on previous work on adrenal function in the viviparous skink *Egernia*  
38 *whitii* (Jones and Bell, 2004; Cartledge et al., 2005), we aim to establish whether  
39 reproductive status modulates adrenocortical activity in females of this species, and to  
40 investigate the endocrine mechanisms that may potentiate differential secretion of  
41 CORT between reproductive stages. We therefore compare the changes in plasma  
42 CORT concentrations in response to the acute stress of capture in females of *Egernia*  
43 *whitii* at three reproductive stages: post-ovulatory, gestating and post-partum. We  
44 hypothesise that both post-ovulatory and gestating (carrying well-developed embryos)  
45 females will down-regulate their acute stress response compared with post-partum  
46 females, due to their current reproductive investment, and that gestating females will  
47 show the most marked reduction. We extend the comparison to males collected at the  
48 same time as the postpartum females, during male reproductive quiescence (Bell,  
49 1997). A comparison of stress responses in males and females when both are  
50 reproductively quiescent is included to determine if there are integral differences in

1 the stress responses of the sexes even when the influence of reproductive steroids  
2 should be minimal.

3  
4 To investigate whether the adrenal response to acute capture stress reflects the  
5 maximal secretory capacity of the adrenal glands in this species, we assess the  
6 adrenocortical response to exogenous ACTH in reproductively quiescent females and  
7 males; however, postovulatory and gestating females are not tested due to concerns  
8 that ACTH stimulation may result in extraordinarily high CORT levels, capable of  
9 disrupting development of embryos. The influence of elevated plasma steroids on  
10 adrenocortical function in female *E. whitii* is investigated by treating reproductively  
11 quiescent females with oestrogen, and measuring plasma concentrations of CORT  
12 before and after injection of ACTH. We chose to use oestradiol -17 because this  
13 steroid is at the end of the steroidogenic pathway, and because the mammalian  
14 literature (see Handa et al., 1994a) provides direct evidence that oestradiol -17  
15 interacts with the HPA axis. We hypothesise that oestrogen -treated females will  
16 exhibit increased basal plasma CORT, and an increased response to exogenous  
17 ACTH.

## 18 19 **2. Methods**

### 20 21 *Animals*

22 Sexually mature (> 65 mm snout-vent length) female and male *Egernia whitii*  
23 were collected from a single site at Orford, Tasmania, Australia (42°34'S, 147°52'E)  
24 during their active season, the austral spring/summer, October through March (1999 –  
25 2000). Females were collected at times of year corresponding to the following stages  
26 of their reproductive cycle: (1) post-ovulation (early-November) (2) gestation (late-  
27 January) and (3) post-partum (early-March): post-partum females were considered to  
28 be reproductively quiescent and the term “gestation” herein refers to females carrying  
29 well formed embryos in the third trimester (Hickman, 1960; Milton, 1987; Chapple,  
30 2005). Males were collected in late March, when they are reproductively quiescent  
31 (Milton, 1987). Males were preliminarily identified by their wider heads and sex was  
32 then confirmed by the presence of eversible hemipenes (Chapple, 2005). Females  
33 were palpated to confirm their reproductive status similar to Chapple (2005) and  
34 Chapple and Keogh (2005). Post-partum females were also quite obvious by their  
35 reduced body weight and flaccid appearance following birth, and follicles were easily  
36 detected in the abdomens of post-ovulatory females.

37  
38 Animals were captured by “fishing” with mealworms and depositing the  
39 animal in a bucket. Blood samples of approximately 100 µL were collected by gently  
40 inserting a heparinised capillary tube into the sub-orbital sinus and allowing the blood  
41 to drip out into a small vial. Blood samples were placed upon ice until return to the  
42 laboratory where plasma was separated from cells by centrifugation and stored at -  
43 20°C until assay for CORT by radioimmunoassay (RIA) (see below).

### 44 45 *Acute Capture Stress*

46 This experiment was conducted in the field. Twenty females were caught at  
47 each reproductive stage and 20 reproductively quiescent males were also captured.  
48 Blood samples were taken immediately after capture for five animals of each group. If  
49 the blood sample was not completed within 5 min of capture, that lizard was not

1 included in the experiment. The remaining animals were held individually in small  
2 cloth bags in a shady position for 10 min, 60 min or 240 min (5 animals per group)  
3 prior to the blood sample being taken. To avoid the influence of possible daily cyclic  
4 variation in plasma CORT concentrations, catching of lizards was timed so that all  
5 post-confinement blood sampling occurred within the same two -hour period (14:00 –  
6 16:00). To avoid recapture of the same individual, no animals were released until all  
7 blood samples had been taken.

#### 8 9 *The ACTH stimulation test*

10 The ACTH stimulation test was conducted in the field in March, using  
11 reproductively quiescent (post-partum) females and males. The aim of this protocol  
12 was to assess the maximum secretory capacity of the adrenal gland. Blood samples  
13 were taken from ten female and ten male *Egernia whitii* immediately upon capture.  
14 Lizards were alternately assigned to ACTH or saline injection as caught. Five lizards  
15 of each sex were injected intramuscularly (forelimb) with 2.5 IU ACTH in 25  $\mu$ L  
16 (Syncanthen Depot, Novartis), while five control lizards of each sex were injected  
17 with 25  $\mu$ L buffered lizard Ringer's solution. All lizards were then held in individual  
18 cloth bags for a period of four hours. This period was chosen because this species'  
19 adrenocortical response to acute stress has been shown to plateau at four hours (Jones  
20 and Bell 2004; this study). At the conclusion of the four -hour period, a second blood  
21 sample was taken, and lizards were released at their point of capture. All blood  
22 samples were placed on ice until return to laboratory where samples were centrifuged  
23 and plasma separated and stored for CORT assay as detailed below.

#### 24 25 *The influence of exogenous oestrogen on adrenal function*

26 Fifteen post-partum females were caught in March. A blood sample was  
27 collected immediately after capture, and the animals were palpated to ensure they  
28 were not still gestating (no gestating females were caught). They were placed into  
29 individual cloth bags for transportation back to the laboratory where they were housed  
30 individually in plastic containers (30 x 20 x 10 cm). Each cage contained a basking  
31 site located under a 25 W light globe on an 13 hr ON: 11 hr OFF cycle. Lizards were  
32 provided with a retreat, and the cage substrate consisted of paper pellets to a depth of  
33 approximately 2 cm. Lizards were fed three times per week on a diet consisting  
34 predominantly of cat food supplemented by tenebrionid larvae (mealworms) and water  
35 was available *ad lib*. The room air temperature was maintained at 20 °C and  
36 fluorescent overhead lighting was set to 13 L: 11 D, mimicking the natural  
37 photoperiod. All females were held under these conditions in the laboratory for 1 wk  
38 prior to the commencement of the experiment, to allow recovery from the increased  
39 adrenal activity associated with the stress of capture and housing. Previous work  
40 indicates that, in this species, 1 wk is sufficient for CORT levels to return to baseline  
41 (Cartledge et al., 2005).

42 After one week of confinement, a blood sample was collected and females  
43 were randomly assigned to one of three treatment groups:

- 44 (i) Saline + ACTH (n=5). Females in this treatment group were administered  
45 50  $\mu$ L of 0.7 % saline intraperitoneally on three consecutive days. On the  
46 fourth day females were administered the ACTH stimulation test as  
47 described in the previous section.
- 48 (ii) Oestrogen + ACTH (n=5). Females in this treatment group received an  
49 intraperitoneal injection of 200 ng oestrogen in 50  $\mu$ L 0.7 % saline on three

1 consecutive days. On the fourth day these animals were administered the  
2 ACTH stimulation test.

- 3 (iii) Oestrogen + Saline (n=4). To control for the likely increase in CORT  
4 concentration due to the stress of injection (as separate from the  
5 stimulatory effects of ACTH injection), females in this treatment group  
6 received an intraperitoneal injection of 200 ng oestrogen in 50  $\mu$ L 0.7 %  
7 saline on three consecutive days. On the fourth day females received an  
8 intramuscular injection of 25  $\mu$ L saline instead of ACTH, and four hours  
9 later a second blood sample was taken.

10  
11 These treatment groups allowed a comparison of plasma CORT concentrations  
12 in low oestrogen (saline injected) against high oestrogen (oestrogen injected) females;  
13 and a comparison of the CORT response s to ACTH in both high oestrogen and low  
14 oestrogen females. In the skink *Tiliqua nigrolutea*, an equivalent dose and injection  
15 regime of oestrogen resulted in elevated plasma oestrogen concentrations after four  
16 days of treatment (Edwards, 1999). Females treated with oestrogen were held in a  
17 separate room to saline treated controls to minimise the possibility of cross  
18 contamination or secondary pheromonal effects due to the oestrogen treatment.

#### 19 *Corticosterone Radioimmunoassay*

20  
21 The radioimmunoassay is described by Jones and Bell (2004). Briefly, CORT  
22 was extracted from plasma aliquots (25 -50  $\mu$ L) with 500  $\mu$ L A. R. grade absolute  
23 ethanol; duplicate aliquots of 200  $\mu$ L of ethanol extract were assayed. Duplicate  
24 standards (0, 25, 50, 100, 200, 400, 800 pg authentic CORT) and sample extracts were  
25 dried down, and 100  $\mu$ L of antiserum (Endocrine Services) and 100  $\mu$ L of tritiated  
26 CORT in phosphate buffered saline were added to each tube. Tubes were incubated at  
27 4°C overnight, and the free and bound fractions separated with dextran-coated  
28 charcoal.

29  
30 Validation of this assay for this species was carried out by Jones and Bell  
31 (2004). The extraction efficiency was 100 % and the amount of hormone that can be  
32 determined as statistically different from the zero tubes was found to be 0.5 ng/mL  
33 plasma. The intra-assay coefficient of variation was 11.2 %, and the inter -assay  
34 variation was 3.4 % (n=5). Serial dilutions of lizard plasma ran parallel to the standard  
35 curve tested over 3-17 ng/mL plasma (Jones and Bell, 2004).

#### 36 *Statistical Analyses*

37  
38 The effects of reproductive stage (females: post -ovulatory, gestating, post-  
39 partum) and time held captive upon plasma CORT concentration during the acute  
40 stress protocol were tested using a factorial design in which females of all  
41 reproductive stages were held captive for all periods of restraint. This design  
42 permitted two-way analysis of variance (ANOVA) within each reproductive state and  
43 between reproductive states for each period of restraint. Variances of mean plasma  
44 CORT concentrations were homogeneous ( $F_{11,44}=0.410, p > 0.5$ ). Following the  
45 ANOVA, a plot of Studentised residuals against predicted values was examined to  
46 check the normality of the residuals. When the ANOVA indicated significant effects,  
47 post-hoc pair-wise comparisons by Tukey's multiple range tests were conducted to  
48 investigate which individual groups differed significantly.

1 Similarly, the responses of reproductively quiescent males were compared to  
2 post-partum females using two-way ANOVA with the factors being sex and time held  
3 captive. Post-hoc pair-wise comparisons were made within each sex using Student -  
4 Neuman-Keuls (SNK) tests. At each time, mean plasma CORT concentrations were  
5 compared between males and females using t-tests.  
6

7 When testing the effect of ACTH injection, the dependent variable analysed  
8 was the change in plasma CORT concentrations pre - and post-injection of ACTH.  
9 Two-way ANOVA was employed to compare plasma CORT concentrations as a  
10 function of the within-groups factor, injection type (experimental or control) and  
11 between-groups factor, sex. When the ANOVA indicated significant effects, post -hoc  
12 pairwise comparisons (SNK tests) were conducted to reveal the points of significant  
13 difference between groups.  
14

15 In the oestrogen injection experiment, repeated -measures analysis of variance  
16 was employed to make comparisons within each treatment group (Saline+ACTH,  
17 Oestrogen+ACTH, Oestrogen+Saline) of CORT concentrations at time zero, 1 wk,  
18 following priming and following ACTH or saline injection. SNK was used for post-  
19 hoc comparisons. To compare between treatment groups at each of these times,  
20 separate one-way ANOVA were employed.  
21

22 Statistical analyses were conducted using the softwares Statistix v1.4  
23 ([www.statistix.com](http://www.statistix.com)) and SAS. In all statistical tests,  $p < 0.05$  was considered  
24 significant.  
25

### 26 **3. Results**

#### 27 *Acute Stress Response*

28 The initial mean plasma CORT concentration did not differ between  
29 reproductive stages ( $F_{2,12}=1.17$ ,  $p > 0.3$ ). However, two-way ANOVA indicated that  
30 the profile of the acute stress response over the 240 min period differed between the  
31 reproductive stages (reproductive stage x time restrained interaction,  $F_{6,60}=2.72$ ,  $p <$   
32  $0.03$ ) (Fig. 1). Post-ovulatory and gestating females exhibited significant elevations in  
33 plasma CORT concentration following both 10 min and 1 h of confinement (post-  
34 ovulatory:  $F_{3,14}=9.17$ ,  $p=0.002$ , gestating:  $F_{3,15}=4.92$   $p < 0.02$ ); however, at 4 h mean  
35 plasma CORT concentrations were no longer significantly different from initial values  
36 in either reproductive state. The mean rise in plasma CORT concentration following  
37 the first 10 min of restraint was significantly greater in post-partum than either post-  
38 ovulatory or gestating females ( $F_{2,12}=8.04$ ,  $p < 0.007$ ). For post-partum females the  
39 mean plasma CORT concentrations at 1 h and 4 h were in fact not significantly below  
40 that of 10 min. Reproductively quiescent males exhibited plasma CORT  
41 concentrations consistently higher than those of post-partum females, but the  
42 difference was significant only at 10 min ( $F_{1,8}=8.89$ ,  $p < 0.02$ ). Concentrations of  
43 plasma CORT varied significantly through time for both sexes ( $F_{3,16}=12.489$ ,  $p <$   
44  $0.001$ ). Reproductively quiescent males exhibited significantly higher plasma CORT  
45 concentrations than at capture at 10 min ( $p < 0.01$ ), 1 h ( $p < 0.01$ ), and 4 h ( $p < 0.01$ )  
46 as compared with time zero.  
47  
48

#### 49 **Figure 1**

50

### 1 *Adrenocorticotrophic hormone (ACTH) stimulation test*

2 Both reproductively quiescent male and female *Egernia whitii* responded to  
3 saline injection with a rise in plasma CORT (Table 1). ACTH injection also induced a  
4 rise in plasma CORT concentration in reproductively quiescent males, but this was not  
5 significantly different from the response to saline injection ( $F_{1,9}=3.2, p > 0.1$ ). In  
6 contrast, the rise in plasma CORT levels exhibited by reproductively active females in  
7 response to ACTH injection was significantly greater than in saline -injected controls  
8 ( $F_{1,8}=10.8, p < 0.05$ ).

### 10 **Table 1**

#### 11 *The influence of exogenous oestrogen on adrenal function*

12 Mean plasma CORT concentrations in females at the four stages of the  
13 oestrogen experiment: immediately following capture, following 1 wk of confinement,  
14 following 4 days of priming with oestrogen or saline, and then lastly on the 12<sup>th</sup> day  
15 either administration of the ACTH stimulation test or otherwise a control saline  
16 injection are shown in Fig. 2. One-way ANOVA revealed that mean plasma CORT  
17 concentrations did not vary significantly between females at time zero ( $F_{2,12}=0.84, p >$   
18  $0.4$ ) or 1 wk ( $F_{2,12}=3.28, p > 0.07$ ), or between females primed with oestrogen or  
19 saline ( $F_{2,12}=3.39, p > 0.07$ ).

### 22 **Figure 2**

23  
24 Comparisons within treatment groups over time revealed no significant  
25 variation in CORT concentration over the course of the experiment for females in the  
26 Oestrogen+Saline treatment group ( $F_{3,12}=1.957, p > 0.17$ ) or the Saline+ACTH  
27 treatment group ( $F_{3,16}=1.819, p > 0.1$ ). However, females in the Oestrogen+ACTH  
28 treatment did display significant variation in CORT between times ( $F_{3,16}=8.712, p <$   
29  $0.01$ ) with concentrations post-ACTH injection higher than pre-injection ( $p < 0.01$ ).

## 32 **4. Discussion**

33  
34 In this study we compared the acute adrenal stress response in female *Egernia*  
35 *whitii* at three reproductive stages: post-ovulation, gestation, and post-partum, to test  
36 the hypothesis that, in this viviparous lizard, gestating females will exhibit a reduced  
37 adrenocortical response compared with post-ovulatory and post-partum females. Basal  
38 plasma CORT concentrations did not differ significantly between post-ovulatory,  
39 gestating and post-partum females in *Egernia whitii*. CORT is a hormone with  
40 catabolic effects associated with increased energy requirements (Greenberg and  
41 Wingfield, 1987; Bentley, 1998). The lack of variation in basal circulating CORT  
42 concentrations in female *Egernia whitii* at different reproductive stages indicates that  
43 females are exhibiting tight regulation over CORT levels even in the presence of  
44 differing energetic demands.

45  
46 However, studies investigating the natural fluctuations in plasma  
47 corticosterone through the annual cycle in female reptiles have generally failed to  
48 demonstrate a consistent pattern of variation with reproductive condition. For  
49 example, in the oviparous lizard *Uta stansburiana* plasma CORT levels are positively  
50 correlated with the relative masses of reproductive tissue (Wilson and Wingfield,

1 1992, 1994), whereas in the parthenogenetic lizard *Cnemidophorus uniparens* plasma  
2 CORT concentrations decline during vitellogenesis and ovulation (Grassman and  
3 Crews, 1990). The oviparous lizard *Cnemidophorus sexlineatus* shows no significant  
4 variation in plasma corticosterone over the reproductive season (Grassman and Hess,  
5 1992b). Similarly, females of the viviparous common gecko *Hoplodactylus maculatus*  
6 show no significant difference in plasma CORT concentrations among four stages of  
7 reproduction (mid/late vitellogenic, mid-pregnant, late pregnant prior to winter  
8 dormancy and spent) (Girling and Cree, 1995). Thus, the way in which plasma CORT  
9 concentration varies with reproductive condition of females shows considerable  
10 variation both within and between viviparous and oviparous lizard species.

11  
12 Reproductively quiescent (post-partum) females exhibited the most marked  
13 immediate response, and the most prolonged response, to acute capture stress. Post -  
14 ovulatory females displayed the most conservative adrenocortical response to acute  
15 capture, with plasma CORT concentrations lower at all times than in the other two  
16 groups of females. In the gestating females, mean CORT concentrations were high (41  
17 ng/mL) at 1 h, similar to those in post-partum females. However, both groups of  
18 reproductive females displayed a recovery by 4 h post-capture, with plasma CORT  
19 concentrations at this time no longer significantly different from initial levels, this was  
20 particularly marked for gestating females. Therefore, although the gestating females  
21 did exhibit initial adrenal responsiveness, their response was attenuated, suggesting  
22 the presence of a mechanism to reduce the duration of the acute stress response. This  
23 may represent an adaptation to mitigate negative impacts of chronically elevated  
24 levels of CORT on gestation: a suite of papers demonstrate that in *Lacerta vivipara*  
25 treating gestating females with exogenous CORT does influence both offspring  
26 phenotype and dispersal behaviour (e.g. Meylan et al. 2002; Belliure et al. 2004;  
27 Meylan and Clobert, 2005). Painter et al. (2002) suggest that placental buffering  
28 means that maternal steroids have little influence on foetal steroid levels in the  
29 viviparous lizard *Sceloporus jarrovi*. The presence of mechanisms for placental  
30 buffering of foetal steroid concentrations may, therefore, in part explain why gestating  
31 females of *E. whitii* exhibited a moderate, though attenuated, stress response, while  
32 post-ovulatory females, in which the chorioallantoic placentae have not yet developed,  
33 exhibited a more marked suppression of the acute stress response. In contrast, in the  
34 viviparous gecko *Hoplodactylus maculatus*, the adrenocortical response to acute  
35 capture in gestating females is as great as that in vitellogenic females (Cree et al.,  
36 2003) and more recent work on gestating females of that species indicates that high  
37 CORT concentrations do not negatively affect the development of the young  
38 following birth (Preest et al., 2005). The authors propose that the capture -induced  
39 CORT levels are lower and of shorter duration than necessary to have deleterious  
40 affects on the embryos and also suggest that the extended period of gestation in this  
41 species may require reduced sensitivity to elevated corticosterone (Preest et al., 2005).  
42 Gravid females of the oviparous *Urosaurus ornatus* show no change in plasma CORT  
43 concentrations in response to an acute stressor, although vitellogenic females respond  
44 with a large increase (Woodley and Moore, 2002). An *in vitro* study of adrenal CORT  
45 production in the presence or absence of ACTH showed no difference between  
46 reproductive stages in the viviparous lizard *Lacerta vivipara* (Dauphin-Villemant et  
47 al., 1990). These results demonstrate that further work is needed on modulation of the  
48 adrenocortical response with reproductive stage in females of oviparous and  
49 viviparous reptiles.

50

1 In comparison with females at all three reproductive stages, reproductively  
2 quiescent males of *Egernia whitii* exhibited the greatest increase in plasma CORT in  
3 response to the acute stressor. Furthermore, CORT concentrations then remained high  
4 over the 4 h period, a pattern similar to that in the post-partum females. This was  
5 unexpected as sex differences in adrenal stress responses have been clearly  
6 demonstrated in mammals, with the majority of work indicating that oestrogen  
7 stimulates CORT production while testosterone ameliorates the acute stress response  
8 (Handa et al., 1994a; Handa et al., 1994b; Viau, 2002; Lund et al., 2004). Whilst very  
9 little is known for reptiles, subordinate male lizards of the species *Anolis*  
10 *carolinensis*, having low plasma testosterone, exhibit a stronger adrenal stress  
11 response than dominant males (Greenberg et al., 1984). As our males were captured  
12 during their period of reproductive quiescence, plasma testosterone concentrations  
13 were low (Bell, 1997), which may explain why their adrenocortical response was  
14 greater than females.

15  
16 Reproductively quiescent males and females responded similarly to exogenous  
17 ACTH. Plasma CORT concentrations increased to 52 ng/mL and 59 ng/mL  
18 respectively, but plasma CORT also increased in the saline-injected animals, reaching  
19 levels of around 30 ng/mL in both sexes, suggesting that the injection induced an acute  
20 stress response. Only in females was the CORT increase caused by ACTH injection  
21 significantly greater than that induced by saline injection alone, partly reflecting the  
22 considerable variation in response between individual males. Carsia and John-Alder  
23 (2003) found that sensitivity to ACTH in isolated adrenal cells varied between males  
24 and females in reproductively quiescent *Sceloporus undulatus*, suggesting a  
25 mechanism for differences in acute stress responses. In *E. whitii*, the plasma CORT  
26 concentrations induced by ACTH injection were comparable with the peak levels  
27 measured during the acute stress protocol. Given the supra-physiological dose of  
28 ACTH employed, this may indicate that this is the maximal concentration of plasma  
29 CORT that can be achieved in this species by stimulation of the HPA axis. Similarly,  
30 Preest et al. (2005) found that exogenous ACTH induced CORT concentrations (62  
31 ng/mL) similar to the maximum (69 ng/mL) measured in pregnant *H. maculatus*  
32 following capture and confinement (Girling and Cree, 1995).

33  
34 To examine the influence of oestrogen upon basal levels of plasma CORT and  
35 the response to ACTH, we treated captive females with oestrogen or saline for four  
36 days, followed by challenge with an injection of ACTH or saline. Plasma CORT  
37 concentrations did not vary between times or between treatments prior to the ACTH  
38 challenge, indicating that exogenous oestrogen did not influence basal CORT  
39 concentrations in this species. This result contrasts with that of Callard and Callard  
40 (1978), who found that oestrogen injection results in an increase in circulating  
41 concentrations of CORT in female *Dipsosaurus dorsalis* in both the reproductively  
42 active and inactive phases of their cycle. Similarly, in mammals, exogenous oestrogen  
43 increases circulating concentrations of both cortisol and corticosteroid-binding  
44 proteins (Coe et al., 1986). However, ovariectomised *Cnemidophorus uniparens*, in  
45 which plasma oestrogen concentrations were significantly decreased, show elevated  
46 levels of CORT in the plasma (Grassman and Crews, 1990), while ovariectomized  
47 *Cnemidophorus sexlineatus* exhibit plasma CORT concentrations that are not  
48 statistically different from those of intact females (Grassman and Hess, 1992a). Such  
49 disparate results suggest that adrenal-gonadal relationships in female reptiles are  
50 complex, reflecting the numerous interactions between CORT, reproductive hormones

1 and metabolic processes (Guillette et al., 1995), and may display a high degree of  
2 inter-species variation.  
3

4 When challenged with ACTH, the oestrogen-treated females responded with a  
5 marked increase in plasma CORT (similar to females subject to the ACTH stimulation  
6 test directly following capture in the field). In contrast, females receiving the saline  
7 control saline injections in place of oestrogen injection showed no such large CORT  
8 response to ACTH injection. This may reflect adrenal exhaustion if these post-partum  
9 females were responding to the acute stress of each saline injection with a large and  
10 extended secretion of CORT, causing exhaustion of adrenal reserves of CORT over  
11 the four days of injection: there is little storage of corticosteroids in the adrenals of the  
12 lizard *Sceloporus undulatus* (Carsia and John-Alder, 2003). However, if the adrenals  
13 were becoming exhausted, we would expect to find undetectable levels of CORT in  
14 these saline-injected females at day four but instead levels were not significantly  
15 different from time zero. Alternatively, saline-injected females may have become  
16 desensitised to ACTH during the four day injection period whilst sensitivity was  
17 maintained in oestrogen-treated females. In mammals, oestrogen can increase cortisol  
18 production by the adrenals (Kitay, 1963), and the response to ACTH is most  
19 pronounced when estrogen levels are highest, such as at pro-oestrous (Viau and  
20 Meaney, 1991). Indeed, in viviparous reptiles, oestrogen is highest during late  
21 vitellogenesis, and drops markedly during early gestation (e.g. Jones and Swain, 1996;  
22 Edwards and Jones, 2001), perhaps suggesting a mechanism for modulation of the  
23 stress response between reproductive phases.  
24

25 In conclusion, our results support our hypothesis that, in comparison with  
26 reproductively quiescent females and males, pregnant females of the viviparous  
27 *Egernia whitii* down-regulate their acute stress response, presumably to protect their  
28 current reproductive investment, with post-ovulatory females showing the most  
29 reduced response. Circulating oestrogen concentrations also appear to modulate the  
30 response to exogenous ACTH, although our results are somewhat equivocal. The  
31 variation in results from the limited number of similar studies in reptiles shows that  
32 further work is required to tease out the mechanisms through which females of  
33 viviparous reptilian species may protect their developing embryos from the potentially  
34 damaging effects of elevated corticosteroids.  
35

### 36 **Acknowledgements**

37  
38 Dr Ashley Edwards, Dr Jane Girling, Natalia Atkins, and Dr Erik Wapstra  
39 kindly assisted with animal capture. Thank you to Dr Stewart Ford for helpful  
40 comments on the manuscript. All procedures were approved by the University of  
41 Tasmania Animal Ethics Committee (Approval no. 99064).  
42

### 43 **References**

44  
45 Astheimer, L. B., Buttemer, W. A. and Wingfield, J. C. (1995). Seasonal and acute  
46 changes in adrenocortical responsiveness in an Arctic-breeding bird. *Horm. Behav.*  
47 29, 442-457.  
48

- 1 Bell, K., 1997. Corticosterone and the stress response in the lizard , *Egernia whitii*  
2 (Scincidae). Unpublished honours thesis, University of Tasmania, Tasmania,  
3 Australia.  
4
- 5 Belliure, J., Meylan, S., Clobert, J., 2004. Prenatal and postnatal effects of  
6 corticosterone on behaviour in juveniles of the common lizard, *Lacerta vivipara*. J.  
7 Exp. Zool. 301A , 401-410.  
8
- 9 Bentley, P.J., 1998. Comparative Vertebrate Endocrinology, 3rd Ed. Cambridge  
10 University Press, Melbourne.  
11
- 12 Burgess, L.H., Handa, R.J., 1992. Chronic estrogen -induced alterations in  
13 adrenocorticotropin and corticosterone secretion, and glucocorticoid receptor -  
14 mediated functions in female rats. Endocrinology 131, 1261-1269.  
15
- 16 Cartledge, V.A., Gartrell, B., Jones, S.M., 2005. Adrenal and white cell count  
17 responses to chronic stress in gestating and postpartum females of the viviparous  
18 skink *Egernia whitii* (Scincidae). Comp. Biochem. Physiol. 141A, 100-107.  
19
- 20 Carsia, R.V., John-Alder, H., 2003. Seasonal alterations in adrenocortical cell function  
21 associated with stress-responsiveness and sex in the eastern fence lizard (*Sceloporus*  
22 *undulatus*). Horm. Behav. 43, 408-420.  
23
- 24 Chapple, D.G. (2005). Life history and reproductive ecology of White's skink,  
25 *Egernia whitii*. Aust. J. Zool. 53, 353-360.  
26
- 27 Chapple, D.G., Keogh, J.S. (2005). Complex mating system and dispersal pattern in a  
28 social lizard, *Egernia whitii*. Mol. Ecol. 14, 1215-1227.  
29
- 30 Cree, A., Tyrrell, C.L., Prest, M.R., Thorburn, D., Guillette Jr., L.J., 2003. Protecting  
31 embryos from stress: corticosterone effects and the corticosterone response to capture  
32 and confinement during pregnancy in a live-bearing lizard (*Hoplodactylus maculatus*).  
33 Gen. Comp. Endocrinol. 134, 316-329.  
34
- 35 Dauphin-Villemant, C., Leboulenger, F., Xavier, F., Vaudry, H., 1990. Adrenal  
36 activity in the female lizard *Lacerta vivipara* Jacquin associated with breeding  
37 activities. Gen. Comp. Endocrinol. 78, 399 -413.  
38
- 39 Edwards, A., 1999. Steroids and reproduction in *Tiliqua nigrolutea*. Unpublished PhD  
40 thesis, University of Tasmania, Tasmania, Australia.  
41
- 42 Edwards, A., Jones, S.M., 2001. Changes in plasma progesterone, estrogen and  
43 testosterone concentrations throughout the reproductive cycle in female viviparous  
44 blue-tongued skinks, *Tiliqua nigrolutea* (Scincidae). Gen. Comp. Endocrinol. 122,  
45 260-269.  
46
- 47 Elsey, R.M., Lance, V.A., Joanen, T., McNease, L., 1991. Acute stress suppresses  
48 plasma estradiol levels in female alligators (*Alligator mississippiensis*). Comp.  
49 Biochem. Physiol. 100A, 649-651.  
50

- 1 Gaskin, J.H., Kitay, J.I., 1970. Adrenocortical function in the hamster. Sex differences  
2 and effects of gonadal hormones. *Endocrinology* 87, 779-786.  
3
- 4 Girling, J.E., Cree, A., 1995. Plasma corticosterone levels are not significantly  
5 related to reproductive stage in female common geckos (*Hoplodactylus maculatus*).  
6 *Gen. Comp. Endocrinol.* 100, 273-281.  
7
- 8 Grassman, M., Crews, D., 1990. Ovarian and adrenal function in the  
9 parthenogenetic whiptail lizard, *Cnemidophorus uniparens* in the field and laboratory.  
10 *Gen. Comp. Endocrinol.* 76, 444-450.  
11
- 12 Grassman, M., Hess, D.L., 1992a. Sex differences in adrenal function in the lizard  
13 *Cnemidophorus sexlineatus*: I. Seasonal variation in the field. *J. Exp. Zool.* 264, 177 -  
14 182.  
15
- 16 Grassman, M., Hess, D.L., 1992b. Sex differences in adrenal function in the lizard  
17 *Cnemidophorus sexlineatus*: II. Responses to acute stress in the laboratory. *J. Exp.*  
18 *Zool.* 266, 183-188.  
19
- 20 Greenberg, N., Wingfield, J.C., 1987. Stress and reproduction: reciprocal  
21 relationships. In: Norris, D.O., Jones, R.E. (Eds.), *Hormones and Reproduction in*  
22 *Fishes, Amphibians, and Reptiles*, 4<sup>th</sup> Ed. Plenum Press, New York, pp 461-503.  
23
- 24 Greenberg, N., Chen, T., Crews, D., 1984. Social status, gonadal state, and the  
25 adrenal stress response in the lizard, *Anolis carolinensis*. *Horm. Behav.* 18, 1-11.  
26
- 27 Guillette Jr, L.J., Cree, A., Rooney, A.A., 1995. Biology of stress: interactions with  
28 reproduction, immunology and intermediary metabolism. In: Warwick, C., Frye, F.L.,  
29 Murphy, J.E. (Eds.), *Health and welfare of captive reptiles*. Chapman & Hall, London,  
30 pp. 32-81.  
31
- 32 Handa, R.J., Burgess, L.H., Kerr, J.E., O'Keefe, J.A., 1994a. Gonadal steroid  
33 hormone receptors and sex differences in the hypothalamo-pituitary-adrenal axis.  
34 *Horm. Behav.* 28, 464-476.  
35
- 36 Handa, R.J., Nunley, K.M., Lorens, S.A., Louie, J.P., McGivern, R.F., Bollnow, M.R.,  
37 1994b. Androgen regulation of adrenocorticotropin and corticosterone secretion in the  
38 male rat following novelty and foot shock stressors. *Physiol. Behav.* 55, 117-124.  
39
- 40 Hansen, D.K., LaBorde, J.B., Wall, K.S., Holson, R.R., Young, J.F., 1999.  
41 Pharmacokinetic considerations of dexamethasone-induced developmental toxicity in  
42 rats. *Toxicol. Sci.* 48, 230-239.  
43
- 44 Hickman, J. L. (1960). Observations on the skink lizard *Egernia whitii* (Lacépède).  
45 *Pap. Proc. Roy. Soc. Tasmania* 94, 111-118.  
46
- 47 Jessop, T.S., 2001. Modulation of the adrenocortical stress response in marine turtles  
48 (Cheloniidae): evidence for a hormonal tactic maximising maternal reproductive  
49 investment. *J. Zool.* 254, 57-65.  
50

- 1 Jones, S.M., Bell, K., 2004. Plasma corticosterone concentrations in males of the  
2 skink *Egernia whitii* during acute and chronic confinement, and over a diel period.  
3 *Comp. Biochem. Physiol.* 137A, 105-113.  
4
- 5 Jones, S.M., Swain, R., 1996. Annual reproductive cycle and annual cycles of  
6 reproductive hormones in plasma of female *Niveoscincus metallicus* (Scincidae) from  
7 Tasmania. *J. Herpetol.* 30, 140-146.  
8
- 9 Kitay, J.I., 1963. Pituitary adrenal function in the rat after gonadectomy and gonadal  
10 hormone replacement. *Endocrinology* 78, 253-260.  
11
- 12 Klose, S. M., Smith, C. L., Denzel, A. J. and Kalko, E. K. V. 2006. Reproduction  
13 elevates the corticosterone stress response in common fruit bats. *J. Comp. Physiol.*  
14 192, 341-350.  
15
- 16 Lance, V.A., Elsey, R.M., 1986. Stress-induced suppression of testosterone secretion  
17 in male alligators. *J. Exp. Zool.* 239, 241-246.  
18
- 19 Lund, T.D., Munson, D.J., Haldy, M.E., Handa, R.J., 2004. Androgen inhibits, while  
20 oestrogen enhances, restraint-induced activation of neuropeptide neurones in the  
21 paraventricular nucleus of the hypothalamus. *J. Neuroendocrinol.* 16, 272-278.  
22
- 23 Mahmoud, I.Y., Guillette Jr, L.J., McAsey, M.E. and Cady, C., 1989. Stress-induced  
24 changes in serum testosterone, estradiol-17 $\beta$  and progesterone in the turtle *Chelydra*  
25 *serpentina*. *Comp. Biochem. Physiol.* 93A, 423-427.  
26
- 27 Meylan, S., Belliure, J., Clobert, J, de Fraipont, M., 2002. Stress and body condition  
28 as prenatal and postnatal determinants of dispersal in the common lizard (*Lacerta*  
29 *vivipara*). *Horm. Behav.* 42, 319-326.  
30
- 31 Meylan, S., Clobert, J., 2005. Is corticosterone-mediated phenotype development  
32 adaptive? Maternal corticosterone treatment enhances survival in male lizards. *Horm.*  
33 *Behav.* 48, 44-52.  
34
- 35 Milton, D.A., 1987. Reproduction in two closely related skinks, *Egernia modesta* and  
36 *E. whitii* (Lacertilia: Scincidae) in south-east Queensland. *Aust. J. Zool.* 35, 35-41.  
37
- 38 Moore, I.L., Jessop, T.S., 2003. Stress, reproduction, and adrenocortical modulation in  
39 amphibians and reptiles. *Horm. Behav.* 43, 39-47.  
40
- 41 Moore, I.T., Greene, M.J., Mason, R.T., 2001. Environmental and seasonal  
42 adaptations of the adrenocortical and gonadal responses to capture stress in two  
43 populations of the male garter snake, *Thamnophis sirtalis*. *J. Exp. Zool.* 289, 99-108.  
44
- 45 Moore, M.C., Thompson, C.W., Marler, C.A., 1991. Reciprocal changes in  
46 corticosterone and testosterone levels following acute and chronic handling stress in  
47 the tree lizard, *Urosaurus ornatus*. *Gen. Comp. Endocrinol.* 81, 217-226.  
48
- 49 Neumann, I.D., Johnstone, H.A., Hatzinger, M., Liebsch, G., Shipston, M., Russell,  
50 J.A., Landgraf, R., Douglas, A.J., 1998. Attenuated neuroendocrine responses to

1 emotional and physical stressors in pregnant rats involve adeno-hypophysial changes.  
2 J. Physiol. 508, 289-300.  
3  
4 Painter, D., Jennings, D.H., Moore, M.C., 2002. Placental buffering of maternal  
5 steroid hormone effects in fetal and yolk hormone levels: a comparative study of a  
6 viviparous lizard, *Sceloporus jarrovi*, and an oviparous lizard, *Sceloporus graciosus*.  
7 Gen. Comp. Endocrinol. 127, 105-116.  
8  
9 Preest, M.R., Cree, A., Tyrrell, C.L., 2005. ACTH-induced stress response during  
10 pregnancy in a viviparous gecko: no observed effect on offspring quality. J. Exp.  
11 Zool. 303A, 823-835.  
12  
13 Valverde, R.A., Owens, D.W., MacKenzie, D.S., Amoss, M.S., 1999. Basal and  
14 stress-induced corticosterone levels in olive ridley sea turtles (*Lepidochelys olivacea*)  
15 in relation to their mass nesting behavior. J. Exp. Zool. 284, 652-662.  
16  
17 Viau, V., 2002. Functional cross-talk between the hypothalamic-pituitary-gonadal and  
18 -adrenal axes. J. Neuroendocrinol. 14, 506-513.  
19  
20 Viau, V., Meaney, M.J., 1991. Variation in the hypothalamic-pituitary-adrenal  
21 response to stress during the estrous cycle in the rat. Endocrinology 129, 2503-2511.  
22  
23 Weinstock, M., 2001. Alterations induced by gestational stress in brain morphology  
24 and behaviour of the offspring. Prog. Neurobiol. 65, 427-451.  
25  
26 Wilson, B.S., Wingfield, J.C., 1992. Correlation between female reproductive  
27 condition and plasma corticosterone in the lizard *Uta stansburiana*. Copeia 3, 691-  
28 697.  
29  
30 Wilson, B.S., Wingfield, J.C., 1994. Seasonal and interpopulation variation in  
31 plasma levels of corticosterone in the side-blotched lizard (*Uta stansburiana*). Physiol.  
32 Zool. 67, 1025-1049.  
33  
34 Wingfield, J. C. 2003. Control of behavioural strategies for capricious environments.  
35 Anim. Behav. 66, 807-816.  
36  
37 Wingfield, J.C., Vleck, C.M., Moore, M.C., 1992. Seasonal changes of the  
38 adrenocortical response to stress in birds of the Sonoran desert. J. Exp. Zool. 264,  
39 419-428.  
40  
41 Wingfield, J.C., Maney, D.L., Breuner, C.W., Jacobs, J.D., Lynn, S., Ramenofsky, M.,  
42 Richardson, R.D., 1995. Modulation of the adrenocortical responses to acute stress in  
43 arctic birds: a possible ecological basis. Am. Zool. 35, 285-294.  
44  
45 Woodley, S.K., Moore, M.C., 2002. Plasma corticosterone response to an acute  
46 stressor varies according to reproductive condition in female tree lizards (*Urosaurus*  
47 *ornatus*). Gen. Comp. Endocrinol. 128, 143-148.  
48

1 *Captions to figures*

2 Fig. 1. Profile of plasma corticosterone concentration (ng/mL) over 4 h of acute  
3 capture stress in *Egernia whitii*. This investigation was carried out in females of three  
4 reproductive states: post-ovulatory, gestating and post-partum and in reproductively  
5 quiescent males. Values are mean  $\pm$  SE. (20 females/males per reproductive state, i.e.  
6 n=5 per experimental group).

7

8

9 Fig. 2. Plasma corticosterone concentrations of post -partum females upon capture in  
10 the field, after 7 d confinement, and following priming with either oestrogen (200  
11 ng/lizard/day) or saline (0.7%) depending on treatment group. Females were primed  
12 with either saline (injection control) or oestrogen and then given either the ACTH  
13 stimulation test or a saline injection (experimental control) . Values are mean  $\pm$  SE. \*  
14 indicates significant difference from 11 d (pre -injection) concentration.


1 Table 1

2 Plasma corticosterone concentrations changes in male and female *Egernia whitii* in response to ACTH  
3 (25IU) versus saline injection. Plasma corticosterone concentrations ( $\text{ng}\cdot\text{ml}^{-1} \pm \text{SE}$ ) pre- and post-  
4 injection are presented along with the difference. Values represent post -injection minus pre-injection.  
5 Five lizards were included in each treatment group.


	Treatment	Plasma Corticosterone Concentration ( $\text{ng}\cdot\text{ml}^{-1} \pm \text{SE}$ )		
		Initial	Post-injection	Difference
<b>Males</b>	Saline control	$10.4 \pm 2.18$	$29.5 \pm 8.06$	$19.1 \pm 9.36$
	ACTH (25 IU)	$12.2 \pm 3.40$	$52.1 \pm 12.6$	$39.9 \pm 6.72$
<b>Females</b>	Saline control	$9.5 \pm 4.41$	$31.1 \pm 11.07$	$21.6 \pm 5.90$
	ACTH (25 IU)	$2.2 \pm 1.78$	$59.4 \pm 17.14$	$57.2 \pm 9.13$ *

6 \*  $p < 0.05$ , significantly different from female controls


7



1



1



1