

# **Environmental influences on reproduction in Antarctic krill, *Euphausia superba***

by Toshihiro Yoshida BE, ME  
Environmental Engineering for Symbiosis  
Soka University, Tokyo, Japan

Submitted in fulfilment of the requirements for the degree of  
Doctor of Philosophy

Institute of Antarctic and Southern Ocean Studies  
University of Tasmania

Oct 2009

## **Declaration**

---

This thesis contains no material which has been accepted for a degree or diploma by any institution. To the best of my knowledge the thesis does not contain any material written or published by another person, except where due reference is made.

Toshihiro Yoshida

## **Authority of access**

---

This thesis is not to be made available for loan or copying for two years following the date this statement was signed. Following that time the thesis may be made available for loan and limited copying in accordance with the Copyright Act 1968.

Toshihiro Yoshida

## Abstract

---

Antarctic krill (*Euphausia superba*) is a keystone species in the Antarctic marine ecosystem, being a major food source for most Antarctic predators. Understanding krill reproduction is important since reproductive output is one of the key determinants of population size. Environmental parameters (food, light and temperature) are generally thought to be responsible for regulating the seasonal maturity cycle of organisms. This study investigates the effect of these environmental parameters on reproduction of krill from several aspects including maturation of female krill, development and hatching of eggs and larval survival.

Female krill were incubated for 10 months under controlled conditions to determine the influence of light and diet on sexual maturity and lipid and fatty acid (FA) compositions. Results from this investigation indicated that endogenous rhythm governed krill sexual maturation. The external sexual characteristics of krill were found to have undergone regression and re-maturation regardless of whether they had been kept under the natural photoperiod (emulating the field environment), or complete darkness. However, photoperiod significantly influenced the ovarian maturation cycle. Diet, either excess or limiting, did not affect this maturation and regression process. There was no significant effect of the addition of carnivorous diet (clam meat) on krill maturity. During five months of low food treatment, lipid was mainly utilized from digestive gland stores in the first two months, and later from other body fractions. Various feeding treatments did not induce any difference in FA compositions in krill. The FA composition of krill was however influenced by sexual maturity status.

After hatching, the larvae of *E. superba* undergo three non-feeding larval stages. This development is dependent on maternal fuel stores laid down in the eggs, which is in turn a function of the dietary conditions prior to spawning. To investigate the effect of maternal diet on embryogenesis, eggs were collected from both laboratory reared krill incubated under different feeding conditions and gravid krill caught in the field. Hatching success, and lipid and FA content and composition differed among the treatments. Positive correlations were found between hatching success and the levels of some specific fatty acids, namely docosahexaenoic acid, linoleic acid and arachidonic acid. Maternal diet to some degree influences the FA composition of eggs, and specific FA may play an important role in embryogenesis of krill eggs.

Krill larvae cope with extremely low phytoplankton concentrations during their first winter, a period thought to be most critical. To investigate the starvation tolerance of larvae, the ultrastructure of digestive gland cells was investigated. The digestive gland is the major site of absorption and storage of nutrients. The effect of starvation on the ultrastructure of the digestive gland of krill larvae was observed as early as 5 days, and these effects increased with duration of starvation. Lipid droplets were rarely found in the digestive cells regardless of whether the larvae were fed or starved. These observations indicate that krill larvae utilize all available nutrients for growing rather than laying down storage. This process indicates that krill larvae are adapted to an environment where the food supply is continuous.

Increasing our understanding of Antarctic krill population is imperative to improve management model for the krill fishery. An understanding of factors influencing fecundity and annual recruitment of krill is important in determining population size. The results of this research helped elucidate the influence of environmental parameters on reproduction of krill. This research also facilitated a better understanding of how krill adapt to the extreme seasonal conditions, which makes them one of the most successful species in Southern Ocean ecosystem.

# Table of Contents

---

Declaration .....	ii
Authority of access .....	ii
Abstract.....	iii
Table of Contents.....	v
Acknowledgements .....	vii
Co-Authorship.....	viii
<b>Chapter 1 General Introduction.....</b>	<b>1</b>
1.1 Introduction .....	1
1.2 Maturation of Antarctic krill.....	1
1.3 Larval ecology.....	3
1.4 Overwintering biology of Antarctic krill.....	4
1.5 The role of lipid in Antarctic krill .....	5
1.6 Aims and hypothesis of thesis .....	6
<b>Chapter 2 Maturation cycle of Antarctic krill: Do diet and food matter?.....</b>	<b>9</b>
2.1 Abstract .....	9
2.2 Introduction.....	9
2.3 Materials and Methods.....	11
2.3.1 Collection of krill.....	11
2.3.2 Aquaria conditions.....	11
2.3.3 Determination of sexual maturity stages.....	13
2.3.4 Measurement of respiration rate.....	13
2.4 Results.....	14
2.4.1 Maturation of thelycum.....	14
2.4.2 Maturation of ovaries.....	15
2.4.3 Respiration rate.....	15
2.4.4 Spawning events.....	17
2.5 Discussion .....	17
2.5.1 Role of light on krill maturation.....	17
2.5.2 Role of diet on krill maturation .....	19
2.5.3 Factors affecting the termination of reproduction of krill.....	19
2.5.4 Respiration rate and maturation of krill .....	21
2.5.5 Effect of experimental condition on krill maturation .....	22
2.5.6 Effect of carnivory on krill maturation.....	22
2.6 Conclusion.....	23
<b>Chapter 3 Dietary influences on the condition of female Antarctic krill during maturation: lipid and fatty acid composition.....</b>	<b>25</b>
3.1 Abstract .....	25
3.2 Introduction.....	26
3.3 Materials and Methods .....	27
3.3.1 Collection of krill.....	27
3.3.2 Aquaria conditions.....	27
3.3.3 Lipid class analysis .....	28
3.3.4 Fatty acid analysis.....	29
3.3.5 Data analysis.....	29
3.4 Results.....	29
3.4.1 Food quality.....	29
3.4.2 Dry mass and total lipid.....	30
3.4.3 Lipid class composition .....	32
3.4.4 Fatty acid composition .....	33
3.4.5 Principal components analysis of FA data .....	33
3.5 Discussion .....	37
3.5.1 Lipid utilization in krill under low food condition .....	37
3.5.2 Variation in fatty acid composition of krill under low food condition.....	40
3.5.3 Differences of lipid class and FA composition in various maturation stages and body fractions of krill .....	40
3.5.4 Fatty acid ratios as dietary markers of carnivory .....	41
3.5.5 Role of omnivory in krill.....	42

3.6 Conclusion.....	43
<b>Chapter 4 Factors determining the hatching success of Antarctic krill embryo: lipid and fatty acid composition.....</b>	<b>45</b>
4.1 Abstract .....	45
4.2 Introduction .....	45
4.3 Materials and Methods .....	47
4.3.1 Embryos from the laboratory.....	47
4.3.2 Food quality of maternal krill in the laboratory .....	47
4.3.3 Embryos from the field .....	47
4.3.4 Determining hatching success .....	48
4.3.5 Lipid analysis.....	48
4.3.6 Data analysis.....	49
4.4 Results.....	49
4.4.1 Food quality.....	49
4.4.2 Hatching success, dry mass and lipid contents of embryos .....	51
4.4.3 Fatty acids.....	51
4.4.4 Temporal differences in lipid and fatty acid profiles.....	51
4.5 Discussion .....	54
4.5.1 Lipid utilization of krill embryos .....	54
4.5.2 Hatching success and FA composition of krill embryos.....	54
4.5.3 TL and FA composition of embryos and food quality of maternal krill .....	55
4.5.4 Hatching success of krill in the field .....	55
4.5.5 Within group hatching success .....	56
4.6 Conclusion.....	57
<b>Chapter 5 Structural changes in the diestive glands of larval Antarctic krill during starvatoin. ....</b>	<b>59</b>
5.1 Abstract .....	59
5.2 Introduction .....	59
5.3 Materials and methods .....	60
5.3.1 Collection of experimental larvae.....	60
5.3.2 Experimental condition.....	61
5.3.3 Transmission electron microscope (TEM) analysis .....	61
5.4 Results.....	61
5.4.1 Fed larvae .....	61
5.4.2 Starved larvae.....	63
5.5 Discussion .....	66
5.5.1 Effect of starvation on R-cell and larval survival .....	66
5.5.2 Lipid droplets in R-cells of larvae .....	67
5.5.3 Ability to take up food after starvation.....	67
5.6 Conclusion.....	67
<b>Chapter 6 General Discussion.....</b>	<b>69</b>
6.1. Effect of light on maturation of female krill .....	69
6.2 Effect of diet on maturation of female krill.....	70
6.3 Endogenous rhythm and maturation of female krill .....	70
6.4 Conceptual model linking environmental parameters and maturation of female krill .....	71
6.5 Lipid transfer from diet - female krill and eggs.....	72
6.6 Embryonic and larval development.....	73
6.7 Conclusion .....	73
References.....	75
Appendices .....	85
Appendix. 1. Kawaguchi S, Yoshida T, Finley L, Cramp P, Nicol S (2007) The krill maturity cycle: a conceptual model of the seasonal cycle in Antarctic kill. <i>Polar Biol</i> 30: 689-698	
Appendix. 2. Hagen W, Yoshida T, Virtue P, Kawaguchi S, Swadling K, Nicol S, Nichols P (2007) Effect of a carnivorous diet on the lipids, fatty acids and condition of Antarctic krill, <i>Euphausia superba</i> . <i>Antarct Sci</i> 19: 183-188	

## Acknowledgements

---

I would like to thank So Kawaguchi for his continuous support and encouragement throughout this research. His academic rigour has instilled discipline in my work. Being from the same culture, his efforts and dedication on his work have been my motivation to overcome all difficulties during my research journey in Australia.

I would like to acknowledge Patti Virtue for her generous support and encouragement, and her expertise in bridging between both ecological and chemical aspects of this research. She showed concern not only in my study but also in my life as an international student. Her warm encouragement instilled confidence in me from time to time.

I would like to thank Peter Nichols for his supervision on the chemical aspect of this research, and his generous help with my work at CSIRO. He has been fair and impartial with my research efforts and offered enthusiastic support with his feedback on my chapter submissions.

Having the above three professionals as my research supervisors was the most fortunate thing during my PhD research. I deeply acknowledge for their generosity and patience.

I would also like to thank George Jackson for various useful comments during early stage of my research.

Steve Nicol is acknowledged for his critique and review of this thesis.

I would like to acknowledge Kelvin Michel for his valued administration, enabling me to complete my PhD as an international student.

I would not have been able to start the whole experiment without the support of Robert King, Paul Cramp and Andrew Mceldowney, the krill team at the Australian Antarctic Division.

I would like to thank Mathias Teschke for his support with the laboratory experiment and sharing beautiful Tasmania with me.

I thank the officers and crew of RV 'Aurora Australis' for their professional and friendly support during the marine research cruise BROKE-WEST.

I would like to thank Matt Miller, Mina Augerinos, Ben Mooney, Peter Mansour and Daniel Holdsworth, for technical assistance with lipid analysis at CSIRO.

Aquarium keepers in the Port of Nagoya Public Aquarium, Yasuo Hirano and Tsuyoshi Matsuda, are acknowledged for enabling me with basic skills in cultivating krill and offering advice during my PhD study.

My enrolment at the University of Tasmania was supported by an International Postgraduate Research Scholarship from the Australian Government. I gratefully acknowledge as well, a partial stipend scholarship from ACE/CEC and Antarctic Tasmania.

To all my friends who drunk with me, encouraged me, and shared a beautiful time with me, thank you.

Last but not least, I express thanks to my loving family. Mum and Dad, your positive attitude and encouragement with conviction have inspired me. Thank you.

お父さん、お母さん、ありがとう。

## Co-Authorship

---

The following people contributed to the publication of the work undertaken as part of this thesis.

Yoshida T, Kawaguchi S, Meyer B, Virtue P, Penschow J, Nash G (2009) Structural changes in the digestive glands of larval Antarctic krill (*Euphausia superba*) during starvation. *Polar Biol* 32(3): 503-507

The following people have contributed to the following chapters.

- So Kawaguchi, Patti Virtue and Peter Nichols assisted with the general supervision of all aspects of this thesis. These included experimental design, interpretation of data and proof reading manuscripts.
- Bettina Meyer assisted with experimental design, interpretation of data and proofreading manuscript in Chapter 5.
- Jennifer Penschow and Gerry Nash assisted technically with TEM, including preparation of samples and operation of TEM in Chapter 5.

We the undersigned agree with the above stated “proportion of work undertaken” for the above published peer-reviewed manuscripts contribution to this thesis.

Supervisor:  
Dr. So Kawaguchi

Supervisor:  
Dr. Patti Virtue

Supervisor  
Dr. Peter Nchols