

THE GEOLOGICAL SETTING AND FORMATION OF  
THE ROSEBERY VOLCANIC-HOSTED MASSIVE  
SULPHIDE OREBODY, TASMANIA.

by Geoffrey R. Green, B.Sc.(Hons.)

A thesis submitted in partial fulfillment of the  
requirements for the degree of Doctor of Philosophy.

UNIVERSITY OF TASMANIA

HOBART

1983

Except as stated herein, this thesis contains no material which has been accepted for the award of any other degree or diploma in any University, and, to the best of my knowledge and belief, this thesis contains no copy or paraphrase of material previously published or written by another person, except where due reference is made in the text of the thesis.

A handwritten signature in cursive script that reads "Geoffrey R. Green". The letters are fluid and connected, with a prominent initial 'G'.

Geoffrey R. Green.

September, 1983

*"Rosebery: A tale of despair."*

*... Blainey (1954)*

## CONTENTS

	page
ABSTRACT	vi
Chapter 1 INTRODUCTION	1.1
Previous literature	1.2
Acknowledgements	1.3
Chapter 2 REGIONAL GEOLOGY	2.1
2.1 Precambrian framework	2.1
2.2 Clastic sequences of the Dundas Trough	2.3
2.3 Mount Read Volcanics	2.8
(i) Central sequence	2.8
(ii) Western sequence	2.10
(iii) Tyndall Group	2.11
2.4 Late Cambrian to Devonian sequences	2.12
2.5 Devonian deformation	2.13
2.6 Devonian to Late Carboniferous granitoid emplacement and mineralization	2.13
Chapter 3 GEOLOGY OF THE ROSEBERY AREA	3.1
3.1 Mount Read Volcanics	3.1
(i) Footwall pyroclastics	3.1
(ii) Host rock	3.6
(iii) Black slate	3.7
(iv) Massive pyroclastics	3.9
(v) Mount Black Volcanics	3.14
(vi) Cauldron subsidence	3.16
3.2 Rosebery Group	3.17
(i) Eastern units	3.19
(a) Chamberlain Shale	3.19
(b) Stitt Quartzite	3.21
(c) Westcott Argillite	3.27
(ii) Central units	3.29
(a) Mudstone-dominated sequence	3.29
(b) Salisbury Conglomerate	3.31
(c) Natone Volcanics	3.34
(iii) Munro Creek Formation	3.35
3.3 Crimson Creek Formation	3.35
3.4 Structure	3.36
3.5 Environment of deposition and provenance of the Rosebery Group	3.49
3.6 Regional extent of the Rosebery Group and correlates	3.53
3.7 Relationship between the Rosebery Group and the Mount Read Volcanics	3.56
3.8 Tectonic History	3.61
(i) Previous Models	3.61
(ii) Constraints on a Tectonic Model	3.62
(iii) Discussion and Tectonic Model	3.67
Chapter 4 THE ROSEBERY OREBODY: FORM, METAL ZONING AND MINERALOGY	4.1
4.1 Form	4.1
4.2 Gross mineralogy and texture	4.5
4.3 Metal zoning	4.8

	page
4.4 Mineralogical zonation	4.18
(i) Arsenopyrite	4.18
(ii) Sphalerite	4.21
(iii) Pyrrhotite	4.25
(iv) Magnetite and hematite	4.26
(v) Barite	4.27
(vi) Albite	4.27
(vii) Chlorite	4.27
(viii) Sericite	4.33
(ix) Carbonate	4.33
Chapter 5 HYDROTHERMAL ALTERATION	5.1
5.1 Geological and textural relations	5.1
5.2 Chemical changes	5.5
5.3 Comparison with other deposits	5.10
5.4 Trace elements in footwall pyrite	5.13
Chapter 6 SULPHUR ISOTOPE INVESTIGATION	6.1
6.1 Introduction	6.1
6.2 Results of the sulphur isotope study	6.3
(i) Sulphide orebody	6.3
(ii) Footwall schist	6.8
(iii) Barite orebody	6.8
6.3 Isotope fractionation between coexisting minerals	6.9
6.4 Source of sulphur	6.12
Chapter 7 METAMORPHISM OF THE OREBODY	7.1
7.1 Introduction	7.1
7.2 Pressure estimates	7.1
(i) Geological estimate	7.1
(ii) Sphalerite geobarometry	7.2
7.3 Pyrite-pyrrhotite sulphur isotope fractionation	7.2
7.4 Biotite and garnet	7.3
7.5 Arsenopyrite-tennantite	7.5
7.6 Chlorite	7.9
7.7 Discussion	7.20
Chapter 8 CONDITIONS OF ORE FORMATION	8.1
8.1 Introduction	8.1
8.2 Chemical environment of ore formation	8.2
8.3 Deposition on the seafloor	8.10
8.4 Generation of the ore solution	8.17
8.5 Features of Rosebery-type deposits	8.20
Chapter 9 CONCLUSIONS	9.1
REFERENCES	R.1
Appendix 1 Specimen catalogue	A1.1
2 Mines Department Drill Hole Rosebery 1	A2.1
3 Pyrite separation method	A3.1
4 Thermodynamic data	A4.1
5 Log ( $\Sigma\text{SO}_4/\Sigma\text{H}_2\text{S}$ ) versus pH and T diagrams	A5.1
6 Sulphur isotope geothermometry	A6.1

## LIST OF TABLES

	page
Table 3.1 Stratigraphy of Rosebery Group.	3.57
4.1 Mineralogy and texture of samples from selected sections.	4.7
4.2 Fit of trend surfaces to metal distribution data.	4.12
4.3 Composition of arsenopyrites.	4.22
4.4 Composition of sphalerites.	4.24
4.5 Composition of gangue minerals.	4.30
5.1 Analyses of altered and unaltered Mt Read Volcanics.	5.7
5.2 Percentage change of components in altered footwall pyroclastics.	5.11
5.3 Co, Ni, and Mn in footwall schist pyrite.	5.14
6.1 Sulphur isotope data.	6.4
6.2 Sulphur isotope fractionation data for metamorphosed massive sulphide deposits.	6.6
7.1 Structural formulae of chlorites.	7.14
7.2 Values of $\log \alpha_{O_2}$ and $X_6$ for Rosebery chlorites	7.15
8.1 Comparison of Rosebery-type and Kuroko-type ores	8.25

## LIST OF FIGURES

Figure 1.1 Locality map of central western Tasmania.	in pocket
2.1 Geological map of western Tasmania.	2.2
2.2 Geological map of central-western Tasmania.	2.4
2.3 Geological history of northwestern Tasmania.	2.6
3.1 Geology of the Rosebery area.	in pocket
3.2 Hand specimen of welded tuff, footwall pyroclastics.	3.2
3.3 Photomicrograph of welded tuff, footwall pyroclastics.	3.2
3.4 Shard texture, footwall pyroclastics.	3.4
3.5 Shard texture, cleaved footwall pyroclastics.	3.4
3.6 Sandstone in black slate.	3.8
3.7 Hand specimen of massive pyroclastics.	3.8
3.8 Ovoids in tuff, massive pyroclastics.	3.13
3.9 Pumice tuff, massive pyroclastics.	3.13
3.10 Welded tuff, north Pieman River area.	3.15
3.11 Breccia, Mt Black Volcanics.	3.15
3.12 Volcaniclastic sandstone, Chamberlain Shale.	3.20
3.13 Amalgamated sandstone beds, Stitt Quartzite.	3.20
3.14 Groove casts, Stitt Quartzite.	3.22
3.15 Granule sandstone, Stitt Quartzite.	3.22
3.16 Sandstone, Stitt Quartzite.	3.24
3.17 B-C turbidite, Stitt Quartzite.	3.24
3.18 Load casted sandstone, Stitt Quartzite.	3.25
3.19 Sandstone bed thickness diagram, Stitt Quartzite.	3.26

List of Figures cont.	page
Figure 3.20 Siltstone, Westcott Argillite.	3.28
3.21 Slide conglomerate, Westcott Argillite.	3.28
3.22 Sphalerite-bearing clast in conglomerate, Westcott Argillite.	3.30
3.23 Geological map of area around fold hinge at 7676E, 7740N.	in pocket
3.24 Graded sandstone bed, Westcott Argillite.	3.30
3.25 Dolerite clast in conglomerate, Westcott Argillite correlate.	3.32
3.26 Salisbury Conglomerate, Pieman River Gorge.	3.32
3.27 Sandstone, Crimson Creek Formation.	3.37
3.28 Fault contact between the Mt Read Volcanics and Stitt Quartzite.	3.39
3.29 Lenticles of sandstone in black slate.	3.39
3.30 E-W cross section through Rosebery Group at 7800mN.	3.41
3.31 Poles to bedding and cleavage, Rosebery Group.	3.43
3.32 E-W cross section through Rosebery at 7350mN.	3.44
3.33 Disrupted bedding in sandstone-mudstone unit.	3.46
3.34 Tectonic breccia in slide zone.	3.46
3.35 Fault contact between Westcott Argillite correlate and Munro Creek Formation.	3.47
3.36 Cross sections through Rosebery Group correlates north of Pieman River.	3.55
3.37 Cross section through correlates of Rosebery Group south of Williamsford.	3.55
3.38 Late Cambrian palaeogeography of Rosebery area.	3.60
3.39 Present tectonic elements of Taupo-Hikurangi zone	3.69
3.40 Evolution of Dundas Trough.	3.74
4.1 Cross section through Rosebery orebody at 500 N.	4.2
4.2 Rosebery mine, geology of 15 level.	4.3
4.3 Rosebery orebody, longitudinal projection.	4.4
4.4 Method of unfolding orebody.	4.10
4.5 Data points, metal zoning study.	4.11
4.6 Third-order trend surfaces for metals -	4.14-
(a) Pb	(b) Zn
(c) Ag	(d) Au
(e) Cu	(f) Fe
(g) Zn/Fe	4.16
4.7 Pre-cleavage arsenopyrite in host rock.	4.19
4.8 Type 1 arsenopyrite.	4.19
4.9 Type 2 arsenopyrite.	4.20

List of Figures continued	Page
4.10 Type 2 arsenopyrite	4.20
4.11 Mole % FeS in Rosebery sphalerites	4.23
4.12 Composition of chlorites from Rosebery	4.29
4.13 Banded sulphide ore	4.31
4.14 Colloform carbonate	4.31
5.1 Blotchy texture of K-feldspar.	5.2
5.2 Quartz-sericite schist.	5.2
5.3 Silicified tuff in massive pyroclastics.	5.6
5.4 Albite-chalcopyrite vein in footwall schist.	5.6
5.5 TiO <sub>2</sub> -Zr plot, Mt Read Volcanics.	5.9
5.6 TiO <sub>2</sub> -Al <sub>2</sub> O <sub>3</sub> plot, Mt Read Volcanics.	5.9
5.7 Co content of footwall schist pyrite.	5.15
6.1 Distribution of sulphur isotope values, sulphide orebody.	6.5
6.2 Sulphur isotope temperatures.	6.10
6.3 Effects of Rayleigh fractionation and disequilibrium sulphide crystallization	6.20
6.4 Change in sulphur source as a function of water: rock ratio	6.32
6.5 Change in $\delta^{34}\text{S}$ values: leaching model	6.33
6.6 Effect of hydrothermal solution: seawater mixing on $\delta^{34}\text{S}$	6.36
7.1 Garnet-chlorite rock.	7.4
7.2 Log $a_{\text{S}_2} \sim 1/T$ diagram showing stability field of arsenopyrite.	7.8
7.3 Log $a_{\text{O}_2} \sim 1/T$ diagram showing conditions of chlorite equilibration, P = saturation pressure of water.	7.16
7.4 Log $a_{\text{O}_2} \sim 1/T$ diagram showing conditions of chlorite equilibration, P = 1 kilobar.	7.17
7.5 Log $a_{\text{O}_2} \sim 1/T$ diagram showing conditions of chlorite equilibration, P = 2 kilobars.	7.18
8.1 Log $\Sigma\text{SO}_4/\Sigma\text{H}_2\text{S} \sim T$ diagram, $m_{\Sigma\text{S}} = 0.002$ .	8.5
8.2 Sketch reconstruction of sulphide orebody.	8.11
8.3 Density-temperature curves for mixing of hydrothermal solutions and seawater.	8.13
8.4 Sketch of possible mode of ore formation from plumes of solution.	8.15
A5.1 Log $\Sigma\text{SO}_4/\Sigma\text{H}_2\text{S} \sim \text{pH}$ diagram 250°C, $m_{\Sigma\text{S}} = 10^{-2}$ .	A5.3
A5.2 Log $\Sigma\text{SO}_4/\Sigma\text{H}_2\text{S} \sim \text{pH}$ diagram 250°C, $m_{\Sigma\text{S}} = 10^{-3}$ .	A5.4
A5.3 Log $\Sigma\text{SO}_4/\Sigma\text{H}_2\text{S} \sim \text{pH}$ diagram 300°C, $m_{\Sigma\text{S}} = 10^{-2}$ .	A5.5
A5.4 Log $\Sigma\text{SO}_4/\Sigma\text{H}_2\text{S} \sim \text{pH}$ diagram 300°C, $m_{\Sigma\text{S}} = 10^{-3}$ .	A5.6
A5.5 Log $\Sigma\text{SO}_4/\Sigma\text{H}_2\text{S} \sim T$ diagram, $m_{\Sigma\text{S}} = 10^{-2}$ .	A5.7