A FIRST LIST OF THE BIRDS OF TASMAN'S PENINSULA.

By J. R. McClymont, M.A.

The subjoined list is not the work of a scientific ornithologist, but of one who has spent a good many hours in ambush, an interested observer of the manners and customs of birds. There are a few of those included in the list which are not of common occurrence in the district above designated; these have been collated with specimens in the Museum, or with Gould's descriptions and plates. The field of observation is within a radius of a mile from the settlement at Koonya, unless otherwise stated in a note. The classification is adopted from "A Systematic List of Tasmanian Birds," by Col. W. V. Legge, R.A., F.L.S.

ACCIPITRES.

- 1. ASTUR NOVÆ HOLLANDIÆ, Gmelin. White Goshawk. Two or three seen within the last four years.
- 2. UROÆTUS AUDAX, Latham. Wedge-tailed Eagle.
- 3. Haliætus leucogaster, *Gmelin*. Grey-backed Sea Eagle. One seen on the Eastern Coast near Tasman's Arch.
- 4. HIERACIDEA ORIENTALIS, Schlegel. Brown Hawk.
- 5. Ninox Boobook, Latham. Brown Hawk Owl. (Ewing in Tas. Jour. i. 53).
- 6. NINOX MACULATA, Vig. and Horsf. Spotted Hawk Owl. Often heard. One was caught in the kitchen of a house, having flown in.

PSITTACI.

- 7. CALYPTORHYNCHUS XANTHONOTUS, Gould. Black Cockatoo. Frequent. Said by the country people to presage rain.
- 8. Platycercus flaviventris, Temminck. Yellow-bellied Parrakeet.
- 9. Pezoporus formosus, Latham. Ground Parrakeet. On the downs of native grass on the N.W. side.
- 10. LATHAMUS DISCOLOR, Shaw. Swift Lorikeet.

PICARIÆ.

- 11. Cuculus Pallidus, Latham. Pallid Cuckoo.
- 12. Cuculus flabelliformis, Latham. Fantailed Cuckoo.
- 12. Podargus cuvieri, Vig. and Horsf. More Pork.

 About the old buildings, from which it is difficult to dislodge them, even with sticks and stones.

PASSERES.

- 14. CORONE AUSTRALIS, Gould. White-eyed Crow.
- 15. STREPERA FULIGINOSA, Gould. Black Magpie. Seen rarely.
- 16. Strepera arguta, Gould. Hill Magpie. As many as twenty together occasionally.
- 17. Grauculus parvirostris, Gould. Summer Bird. Arrives in the latter part of October.
- 18. COLLYRIOCINCLA RECTIROSTRIS, Jard and Selby. Whistling Shrike Thrush.

Known as "Whistling Dick," or locally and anthropomorphically, "Black-eyed Susan." Man's nearness to the animal is rudely indicated in these friendly and familiar appellations.

19. GYMNORHINA HYPERLEUCA, Gould. Magpie.

First domiciled at Saltwater River, this bird has reached Impression Bay. The Sparrow has advanced in the same direction.

- 20. Cracticus cinereus, Gould. Butcher Bird. Its piping note is heard from time to time.
- 21. Pachycephala olivacea, Vig. and Horsf. Olivaceous Thickhead.
- 22. Pachycephala glaucura, Gould. Grey-tailed Thickhead. This bird, or P. gutturalis, was observed once in the gully behind Koonya.
- 23. Rhipidura saturata, Sharpe. Fantail. Called by boys "Cranky Fan" from its erratic movements in pursuit of insects.
- 24. Myiagra Nitida. Gould. Shining Flycatcher. Seen once near Taranna.
- 25. Petræca Leggii, Sharpe. Scarlet-breasted Robin.
- 26. Petræca phænicea, Gould. Flame-breasted Robin. The female has the under surface grey. See Gould i. 283.
- 27. Petræca rhodinogaster, Drapier. Pink-breasted Robin. The nest is exquisitely adorned with lichen.
- 28. Petræca vittata, Quoy et Gaimard. Dusky Robin.
- 29. Malurus Gouldi, Sharpe. Blue Wren.

The males through the winter in blue plumage are seen. See Gould i. 321.

- 30. GEOCICHLA MACRORHYNCHA, Gould. Ground Thrush.
- 31. CINCLOSOMA PUNCTATUM, Lath. Ground Bird.
- 32. Calamanthus fuliginosus, Vig. and Horsf. Rush Warbler.

- 33. STIPITURUS MALACHURUS, Shaw. Emu Wren.
 Seen many years ago at Eagle Hawk Neck; lately, near
 Roaring Beach.
- 34. Sericornis humilis, Gould. Sombre Bush Tit.
- 35. ACANTHIZA DIEMENENSIS, Gould. Brown Tail.
- 36. Acanthorynchus tennirostris, Lath. Spine Bill. Very easily beguiled by an imitation of its whistling notes.
- 37. Melithreptus validirostris, Gould. Strong-billed Honey-eater.
- 38. Melithreptus melanocephalus, Gould. Black-headed Honey-eater.
- 39. PTILOTIS FLAVIGULA, Gould. Yellow-throated Honeyeater.

The Spring note of this bird is almost as characteristic of the season as the note of the Summer-bird.

- 40. Meliornis Novæ Hollandiæ, Lath. New Holland Honey-eater.
- 41. Meliornis Australasiana, Shaw. Tasmanian Honeveater.

The peculiar humming and clicking noise made by this bird is, I suppose, produced by the wings and mandibles.

- 42. ACANTHOCHÆRA INAURIS, Gould. Wattle Bird.
- 43. Acanthochæra Mellivora, Lath. Brush Wattle Bird.
- 44. Zosterops cærulescens, Lath. White Eye.
- 45. PARDALOTUS PUNCTATUS, Shaw and Nodder. Diamond Bird.
- 46. PARDALOTUS AFFINIS, Gould. Allied Diamond Bird.
- 47. HIRUNDO NEOXENA, Gould. Australian Swallow.
- 48. ESTRELDA BELLA, Lath. Fire-tailed Finch.
- 49. Anthus Australis, Vig. and Horsf. Australian Pipit.

COLUMBÆ.

50. Phaps chalcoptera, Latham. Bronze-wing.

GALLINÆ.

A Quail, but species not determined.

FULICARIÆ.

51. TRIBONYX GOULDI, Sclater. Native Hen. One shot near Koonya.

GRALLÆ.

52. ÆGIALITES MONACHA, Geoffrey. Hooded Sand Plover.

53. ÆGIALITES BUFICAPILLA, Temm. Red-capped Sand Plover.

Both found on the Beach of Pirates Bay.

- 54. LOBIVANELLUS LOBATUS, Latham. Wattled Plover. A bird answering to this shot near Safety Cove.
- 55. Hæmatopus Longirostris, Vieill. White-breasted Oyster Catcher.
- 56. Hæmatopus unicolor, Wagler. Sooty Oyster Catcher. GAVIÆ.
- 57. Sterna Caspia, *Pallas*. Caspian Tern. Shot near Taranna.
- 58. STERNA POLIOCERCA, Gould. Bass's Straits Tern.
- 59. LARUS PACIFICUS, Latham. Pacific Gull.
- 60. LARUS NOVÆ HOLLANDIÆ, Stephens. Little Gull.

STEGANOPODES.

- 61. Sula serrator, Banks. Australian Gannet.

 This bird enters the water sidewise, and penetrating it first with one outstretched wing raises a small fountain of foam.
- 62. PHALACROCORAX CARBO, Linu. Common Cormorant.
- 63. Phalacrocorax leucogaster, Gould. White-breasted Cormorant.

ANSERES.

64. Cygnus atratus, Lath. Black Swan.

Seen near Price's Flats. Also on lagoon near Lime Bay, where the nest has been found.

65. Anas superciliosa, *Gmelin*. Wild Duck. On the agoons.

- 66. Anas castanea, Egton. Teal. Shot near Koonya.
- 67. BIZIURA LOBATA, Shaw. Musk Duck. Shot at Impression Bay.

HERODIONES.

- 68. Ardea Novæ Hollandiæ, Lath. White-fronted Heron. Sometimes seen frequenting cultivated paddocks in winter.
- 69. Botaurus Poicilopterus, Wagler. Australian Bittern. Found on the N.W. side.

IMPENNES.

70. EUDYPTULA MINOR, Forster. Little Penguin.

The following species probably belong to the district, but not with sufficient certainty to be placed in the above list:—
Strix castanops, Cuculus plagosus, Phaps elegans, Porphyrio cyanocephalus, Gallinago Australis.