


# **The effects of oxygen and temperature on the physiology of hatching stage Atlantic salmon**



**Elias T. Polymeropoulos**

(Diploma in Biology)

Submitted in fulfilment of the requirements for the degree of Doctor of  
Philosophy (PhD)

University of Tasmania

July 2013

"doing what little one can to increase the general stock of knowledge is as respectable an object of life, as one can in any likelihood pursue"

*(Charles Darwin)*

## Acknowledgements

First and foremost I would like to thank my family for giving me the special opportunity for this Tasmanian journey. Thank you for showing understanding and patience to travel to the furthestmost place on the planet. I would not have been able to do this without your unconditional love and support.

Thanks to Martin Jastroch for paving the academic path to Australia and for never refusing to help and support me in times of need.

I would like to thank Frapps for convincing me to come to Tasmania, his contribution to this thesis despite god knows how many jobs to deal with, his companionship and hospitality (thanks Deirdre!) along this journey. Thanks for teaching me critical thinking, questioning, inventiveness, “accuracy” and persistence at work, and for fun, beer, parmis, laughter and taking the piss at every opportunity.

Thanks also to my supervisor Nick Elliott, who was always there when needed, his constructive criticism and support as well as organising ability.

Special thanks to Brian Taylor for being a genius and being able to fix and salvage anything I have broken over the years!

Thanks to Suzie Currie and Sacha Leblanc for a great time in tropical Sackville and Prince Edward Island. It was a great experience! Additional thanks to Debbie Plouffe and Aqua Bounty Technologies for giving me the opportunity to work on the transgenic salmon.

I would like to thank Rachel Fleming, Emma Crow and Tina Stephens for saving my sanity and keeping me out of trouble in times of Frapps and lots yummy coffee.

Thanks also to the CSIRO crew and the CSIRO ICT centre, in particular Sarah, Andrea and Harry for proof reading this thesis, Richard, Brad, Ben and Peter for showing interest in my research and many constructive discussions.

Thanks to the School of Zoology, especially Stewart Nicol for good inspirations, Richard Holmes and Tony Sprent for great technical support.

Many thanks to all the people I have met along the way who made this journey the greatest experience of my life so far: Martin WP, Ryan B, JD, Cozzy, Pagey, Danish dynamite, Chewy, Malou, CBS, Pulpo, Elsa, Hubert, Martin & Lisa, Mr. Pink, Auntie Margaret, Cherry Pie, Rebecca, Knopwoods Retreat, Joga Bonito FC, The brothers from the poolside and last but not least TUDC.

### **Declaration of Originality**

"This thesis contains no material which has been accepted for a degree or diploma by the University or any other institution, except by way of background information and duly acknowledged in the thesis, and to the best of my knowledge and belief no material previously published or written by another person except where due acknowledgement is made in the text of the thesis, nor does the thesis contain any material that infringes copyright."

### **Authority of Access**

This thesis is not to be made available for loan or copying for two years following the date this statement was signed. Following that time the thesis may be made available for loan and limited copying and communication in accordance with the Copyright Act 1968.

### **Statement of Co-Authorship**

The following people and institutions contributed to the publication of work undertaken as part of this thesis:

*Candidate* = Elias Polymeropoulos (School of Zoology)

*Author 1* = Prof. Peter Frappell (University of Tasmania)

*Author 2* = Dr. Nick Elliott (Commonwealth Scientific and Industrial Research Organisation)

*Author 3* = Dr. Simon Wotherspoon (Institute for Marine and Antarctic Studies)

**Author details and their roles:**

***Respirometry: Correcting for diffusion and validating the use of plastic multiwell plates with integrated optodes.***

*Located in Chapter 2.3. Correcting for diffusion and validating the use of plastic multiwell plates with integrated optodes*

*Candidate was the primary author and author 1 contributed to the idea, its formalisation and development*

*Author 2 and author 3 assisted with refinement and presentation*

*Author 3 derived the mathematical model that enables general correction of oxygen diffusion occurring across polystyrene multiwell plates.*

We the undersigned agree with the above stated “proportion of work undertaken” for each of the above published (or submitted) peer-reviewed manuscripts contributing to this thesis:

Signed: \_\_\_\_\_

*(Name)*

*Supervisor*

*Pro Vice Chancellor (Global engagement)*

University of Tasmania

*(Name)*

*Head of School*

*School of Zoology*

University of Tasmania

Date: \_\_\_\_\_

**Statement of Ethical Conduct (where applicable)**

“The research associated with this thesis abides by the international and Australian codes on human and animal experimentation, the guidelines by the Australian Government's Office of the Gene Technology Regulator and the rulings of the Safety, Ethics and Institutional Biosafety Committees of the University.”

Cover image: Atlantic salmon (*S. salar*) yolk sac alevin.



## Table of Contents

<b>Acknowledgements</b> .....	iii
<b>Table of Contents</b> .....	vi
<b>List of Figures</b> .....	x
<b>List of Tables</b> .....	xii
<b>Abbreviations</b> .....	xiii
<b>Abstract</b> .....	xiv

### **1. General introduction: Effects of environmental and inherent factors on metabolic, cardiorespiratory function and heat shock protein expression in fish** ..... 17

1.1. Thermal tolerance and oxygen limitation .....	17
1.2. Gas exchange in water breathing vertebrates.....	18
1.3. Hypoxia and temperature .....	21
1.4. Generalised physiological response to hypoxia in adult fish.....	22
1.5. Physiological response to hypoxia during early development in fish.....	24
1.6. Temperature and teratogenic factors during fish incubation .....	26
1.7. Study-specific background.....	27
1.8. Objectives and hypotheses. ....	31

### **2. Methods: Measurement of metabolic rate, heart rate, ventilation rate and heat shock protein expression** ..... 35

2.1. Egg, geometry and embryonic and yolk-sac alevin mass and morphometry..	35
2.2. Measurement of metabolic rate in eggs and alevins .....	36
2.3. Correcting for diffusion and validating the use of plastic multiwell plates with integrated optodes .....	37
2.3.1. <i>The polystyrene-based multiwell aquatic respirometer</i> .....	39
2.4. Simultaneous measurement of metabolic rate, heart rate and ventilation rate in individual alevins .....	46
2.5. Validation of the intermittantly sealed flow-through respirometry system ....	49

2.6. Determination of $P_{\text{crit-hatch}}$ in eggs .....	51
2.7. Measurement of perivitelline $PO_2$ ( $PO_{2(\text{in})}$ ) in eggs .....	52
2.8. Calculation of egg capsule oxygen conductance $GO_2$ .....	53
2.9. Heat shock protein (Hsp) immunodetection .....	54

**3. The maternal effect of differences in egg size influence metabolic rate and hypoxia induced hatching in Atlantic salmon (*Salmo salar*) eggs: implications for respiratory gas exchange across the egg capsule .....**

3.1. Abstract .....	57
3.2. Introduction.....	59
3.3. Experimental protocol.....	63
3.3.1. <i>Fish husbandry and maintenance</i> .....	63
3.3.2. <i>Treatments and measurements</i> .....	63
3.3.3. <i>Statistical analysis</i> .....	63
3.4. Results.....	65
3.4.1. <i>Egg geometry and embryo and alevin morphometry</i> .....	65
3.4.2. <i>Mass-specific metabolic rate and hatching</i> .....	66
3.4.3. <i>Perivitelline <math>PO_2</math> (<math>PO_{2(\text{in})}</math>) and <math>GO_2</math></i> .....	68
3.5. Discussion .....	72
3.5.1. <i>Mass-specific metabolic rate and hatching</i> .....	72
3.5.2. <i>Perivitelline <math>PO_2</math> (<math>PO_{2(\text{in})}</math>) and oxygen exchange</i> .....	76

**4. Hypoxic acclimation leads to metabolic compensation after re-exposure to normoxia and increased cardiorespiratory sensitivity to acute hypoxia in Atlantic salmon (*Salmo salar*) yolk-sac alevins .....**

4.1. Abstract .....	83
4.2. Introduction.....	85
4.3 Experimental protocol.....	91
4.3.1. <i>Fish husbandry and maintenance</i> .....	91
4.3.2. <i>Treatments and measurements</i> .....	91
4.3.3. <i>Statistical analysis</i> .....	93
4.4. Results.....	94
4.4.1. <i>Body mass and morphometry</i> .....	94

4.4.2. <i>Mass-specific metabolic rate</i> .....	94
4.4.3. <i>Simultaneous measurement of mass-specific metabolic rate, heart- and ventilation rate</i> .....	97
4.5. Discussion .....	101
4.5.1. <i>Body mass and morphometry</i> .....	101
4.5.2. <i>Mass-specific metabolic rate</i> .....	101
4.5.3. <i>Simultaneous measurement of mass-specific metabolic rate, heart rate and ventilation rate</i> .....	106
<b>5. Acute but not chronic hyperoxia increases metabolic rate and does not alter the cardiorespiratory response in Atlantic salmon (<i>Salmo salar</i>) yolk-sac alevins at 4°C and 8°C</b> .....	<b>111</b>
5.1. Abstract .....	111
5.2. Introduction .....	113
5.3. Experimental protocol .....	117
5.3.1. <i>Fish husbandry and maintenance</i> .....	117
5.3.2. <i>Treatments and Measurements</i> .....	117
5.3.3. <i>Statistical analysis</i> .....	118
5.4. Results .....	119
5.4.1. <i>Body mass and morphometry</i> .....	119
5.4.2. <i>Mass-specific metabolic rate</i> .....	119
5.4.3. <i>Simultaneous measurement of mass-specific metabolic rate, heart- and ventilation rate</i> .....	120
5.5. Discussion .....	123
5.5.1. <i>Body mass and morphometry</i> .....	123
5.5.2. <i>Mass-specific metabolic rate, heart rate and ventilation rate</i> .....	123
<b>6. Growth hormone transgenesis and polyploidy increase metabolic rate, alter the cardiorespiratory response and influence Hsp expression to acute hypoxia in Atlantic salmon (<i>Salmo salar</i>) yolk-sac alevins</b> .....	<b>127</b>
6.1. Abstract .....	127
6.2. Introduction .....	129
6.3. Materials and Methods .....	134
6.3.1. <i>Experimental animals</i> .....	134
6.3.2. <i>Treatments and Measurements</i> .....	134

6.3.3. <i>Statistical analysis</i> .....	135
6.4. Results.....	137
6.4.1. <i>Body mass and morphometry</i> .....	137
6.4.2. <i>Series I: Mass-specific metabolic rate</i> .....	137
6.4.3. <i>Series II: Simultaneous measurement of mass-specific metabolic rate, heart- and ventilation rate</i> .....	138
6.4.4. <i>Series III: Hsp expression</i> .....	141
6.5. Discussion.....	143
6.5.1. <i>Series I and Series II</i> .....	134
6.5.2. <i>Series III:</i> .....	138
<b>7. General discussion</b> .....	<b>151</b>
<b>8. References</b> .....	<b>167</b>
<b>Appendices</b> .....	<b>187</b>

## List of Figures

<b>Figure 1.1.</b> Temperature effects on aquatic animals. ....	18
<b>Figure 1.2.</b> A simplified schematic of the oxygen cascade in fish.....	19
<b>Figure 1.3.</b> Simplified concept of O <sub>2</sub> convection as described by Fick. ....	20
<b>Figure 2.1.</b> <i>S.salar</i> egg under a dissecting microscope.....	35
<b>Figure 2.2.</b> A polystyrene-based well used for aquatic respirometry.. ....	39
<b>Figure 2.3.</b> Temperature dependent O <sub>2</sub> diffusion through a polystyrene-based well and custom built aluminium (non-diffusible) well over time .....	42
<b>Figure 2.4.</b> LogP <sub>wt</sub> -P <sub>a</sub> against time for a polystyrene-based well at select temperatures and for a custom built aluminium well (non-diffusible).. ....	43
<b>Figure 2.5.</b> Change in PO <sub>2</sub> over time in both aluminium non-diffusible and polystyrene-based wells that hold either Alzet micro-osmotic pumps at 37°C containing a 0.1 M sodium sulphite (O <sub>2</sub> scavenging) solution or <i>S. salar</i> eggs at 8°C.. ....	45
<b>Figure 2.6.</b> Calculated rates of O <sub>2</sub> consumption (R) for either Alzet micro-osmotic pumps at 37°C containing a 0.1 M sodium sulphite (O <sub>2</sub> scavenging) solution or <i>S. salar</i> eggs at 8°C.....	46
<b>Figure 2.7.</b> Diagram of the respirometry setup in combination with the optophysiological system to simultaneously measure $\dot{M}O_2$ , $f_H$ and $f_V$ in individual <i>S.salar</i> alevins at different levels of PO <sub>2</sub> .....	47
<b>Figure 2.8.</b> Example of the PO <sub>2</sub> profile of outflowing water leaving the respirometry chamber against time that occurs when the chamber, previously sealed and in which O <sub>2</sub> is depleted by the oxygen consumption of the animal inside, is flushed with fresh incoming water.....	50
<b>Figure 2.9.</b> Comparison between theoretical and calculated $\dot{M}O_2$ when injecting anoxic water into the respirometer.....	51
<b>Figure 3.1.</b> Relationship between egg capsule thickness (mm) and egg volume (mm <sup>3</sup> ) of eggs from maiden spawners .....	65
<b>Figure 3.2.</b> Mass-specific metabolic rate ( $\dot{M}O_2$ ) of mature <i>S. salar</i> eggs, eggs during hatching and alevins from maiden and repeat spawners in normoxia and after progressive, acute hypoxia using closed system respirometry .....	67
<b>Figure 3.3.</b> Perivitteline PO <sub>2</sub> (PO <sub>2(in)</sub> ) for eggs from maiden spawners .....	69
<b>Figure 3.4.</b> Theoretical model of changes in perivitelline PO <sub>2</sub> (PO <sub>2(in)</sub> ) in maiden and repeat <i>S. salar</i> eggs with increasing hypoxia .....	78

<b>Figure 4.1.</b> Schematic diagram of the experimental groups. Fertilised eggs were incubated at 4°C and 8°C in normoxia .....	93
<b>Figure 4.2.</b> Changes in mean $\pm$ S.E. mass-specific $\dot{M}O_2$ in response to a progressive decline in $PO_2$ at 4, 8 and 12°C of <i>S. salar</i> alevins incubated at a) 8°C and b) 4°C with acclimation to normoxic or hypoxic conditions .....	95
<b>Figure 4.3.</b> Mass-specific $\dot{M}O_2$ at different ambient temperatures (4, 8, 12°C) of yolk-sac alevins measured at their acclimation $PO_2$ , either normoxia (21 kPa) or hypoxia (10.5 kPa) for each incubation temperature of a) 8°C and b) 4°C.....	97
<b>Figure 4.4.</b> Changes in a) and b) mass-specific $\dot{M}O_2$ , c) and d) heart rate and e) and f) ventilation rate of yolk-sac alevins acclimated to normoxic and hypoxic ( $PO_2 = 10.5$ kPa, 15 days) conditions, incubated at 8°C and 4°C and measured at 4°C and 8°C ( $T_a$ ) in response to a stepwise, acute hypoxic challenge ( $PO_2 = 21 \rightarrow 15 \rightarrow 10 \rightarrow 5$ kPa).. .....	98
<b>Figure 5.1.</b> Changes in response to progressive, acute hypoxia ( $21 \rightarrow 5$ kPa) in mass-specific $\dot{M}O_2$ of <i>S. salar</i> yolk-sac alevins acclimated to normoxic ( $PO_2 = 21$ kPa) and hyperoxic ( $PO_2 = 28$ kPa) conditions and measured at 4°C and 8°C using closed system respirometry .....	120
<b>Figure 5.2.</b> Changes in response to two stepwise challenges in a) mass-specific $\dot{M}O_2$ , b) $f_H$ and c) $f_V$ of yolk-sac alevins acclimated to normoxic and hyperoxic (hyper; $PO_2 = 28$ kPa, 15 days) conditions and measured at 4°C and 8°C ( $T_a$ )..	122
<b>Figure 6.1.</b> Changes in $\dot{M}O_2$ in response to progressive, acute hypoxia ( $21 \rightarrow 5$ kPa) of 2nNt, 2nTx, 3nNt and 3nTx <i>S. salar</i> yolk-sac alevins measured at 8°C. ....	138
<b>Figure 6.2.</b> Changes in $\dot{M}O_2$ (a), heart rate $f_H$ (b) and ventilation rate $f_V$ (c) in 2nNt and 3nTx <i>S. salar</i> yolk-sac alevins in response to a stepwise ( $PO_2$ of 21, 13 and 5 kPa) challenge of hypoxia .....	140
<b>Figure 6.3.</b> Representative fluorograms of a) Hsp70 and b) Hsc70 c) Hsp90 synthesis in <i>S. salar</i> yolk-sac alevin whole body homogenates.. .....	141
<b>Figure 7.1.</b> Conceptual diagram of the effects of acute and chronic changes in environmental $O_2$ levels on metabolic rate in Atlantic salmon yolk-sac alevins relative to normoxia .....	158
<b>Figure 7.2.</b> Simplified diagram of the concept of metabolic compensation at different measurement temperatures in Atlantic salmon yolk-sac alevins. ....	160

## List of Tables

- Table 3.1.** Linear regression equations of the form  $y = mx + b$  for  $PO_{2(out)}$  against egg volume at different  $PO_{2(out)}$  .....70
- Table 3.2.** Comparative egg geometry alevin mass (wet) and metabolic parameters relevant to  $O_2$  exchange across the egg capsule of eggs and alevins from maiden and repeat spawning *S. Salar*.....71
- Table 4.1.** Morphometry and body mass (wet) parameters of *S. salar* alevins incubated at 4 and 8°C with acclimation to normoxic or hypoxic conditions .....94
- Table 5.1.** Comparisons of morphometric and body mass (wet) parameters of yolk-sac alevins (*S. salar*) incubated at 4°C in normoxia and after chronic hyperoxia ( $PO_2= 28$  kPa; 15 days) exposure ..... 119
- Table 6.1.** Comparisons of morphometric parameters of newly hatched yolk-sac alevins of different genotypes (2nNt, 2nTx, 3nNt, 3nTx) raised at 8°C .....137

## Abbreviations

2n	=	diploid
3n	=	triploid
bpm	=	beats per minute
$\beta\text{O}_2$	=	solubility of oxygen
$f_{\text{H}}$	=	heart rate
$f_{\text{V}}$	=	ventilation rate
GH	=	growth hormone
$\text{GO}_2$	=	oxygen conductance
Hb	=	hemoglobin
Hsp	=	heat shock protein
$\text{KO}_2$	=	Krogh's diffusion coefficient
kPa	=	kilopascal
$\dot{\text{M}}\text{O}_2$	=	metabolic rate
$\text{PO}_2$	=	partial pressure of oxygen
$T_{\text{a}}$	=	ambient temperature
Tx	=	transgenic
Nt	=	non transgenic

## Abstract

Environmental change and its impact on the form, function and adaptive responses of fish, especially in relation to maternal effects or individual genotype, is of major interest to biologists. Oxygen levels and temperature have profound effects on the physiology of fish, and early developmental stages are particularly susceptible to environmental challenges. In this thesis the metabolic, cardiorespiratory and cellular stress responses to acute or chronic changes of these variables in Atlantic salmon (*Salmo salar*) embryos and yolk-sac alevins were investigated.

Eggs and alevins matched their metabolic demand to acute changes in oxygen levels without altering cardiorespiratory function unless oxygen reached critically low levels (5 kPa). In contrast, chronic hypoxia (10.5 kPa for 15 days), but not hyperoxia (28 kPa for 15 days) resulted in functional and structural modifications that enabled metabolic rate to return to the normoxic (21 kPa) pre-exposure level while maintained in hypoxia (10.5 kPa); this indicating a metabolic compensation. On return to acute normoxia, irrespective of the measurement temperature (4, 8, 12°C), metabolic rate was elevated above the pre-hypoxic value observed in normoxia, presumably a result of the structural and/or functional modifications that occurred in hypoxia.

In addition, differences in metabolism and responses to hypoxia were influenced by maternal factors such as egg size, which in turn are determined by maternal body size. Egg size-dependent differences in metabolic rate of embryos were

present and larger embryos from repeat spawning females exhibited increased tolerance to hypoxia, as represented by lower critical oxygen levels for hatching than observed in embryos from smaller eggs from maiden spawners (13 kPa vs. 17 kPa). This result suggests an advantage in embryos from larger eggs whereby the embryo can obtain sufficient oxygen due to the larger surface area of the egg in respect to the embryo's metabolic rate.

Genetic modification for growth (growth hormone transgenesis) or triploidy (three sets of paired chromosomes) increased metabolic rate above a diploid conspecific (both by 8%). This effect was additive in triploid transgenic alevins that also displayed an altered cardiorespiratory response to acute, severe hypoxia (5 kPa). In addition, acute hypoxia did not elicit a cellular stress response, but was associated with differential (reduced) expression of cellular stress proteins (heat shock proteins) that in one case (Hsp90) was dependent on the growth hormone transgenic genotype.

Taken together, the above observations demonstrate that extrinsic as well as intrinsic factors have substantial effects on the physiology of developing fish, and that the overall response to extrinsic factors contains a temporal component. Given the associated effects are reflected in changes in metabolism, it is likely that the effects will significantly impact growth, survival and performance of the developing fish.