

New Australian thraustochytrids: A Renewable Source of Biofuels, Omega-3 Oils and other Bioproducts

by

Kim Jye Lee Chang B. Biotech. (Hons)
(University of Tasmania)

Submitted in fulfilment of
the requirements of the degree of
Doctor of Philosophy
University of Tasmania, August 2013

Declaration

This thesis contains no material that has been accepted for the award of any other degree or diploma in any tertiary institution. To the best of my knowledge this thesis does not contain material written or published by another person, except where due reference is made.

This thesis is not to be made available for loan or copying for two years following the date this statement was signed. The publishers of the papers comprising Chapters 2, 3 and 5 hold the copyright for that content, and access to the material should be sought from the respective journals. Following that time the remaining non published content of the thesis may be made available for loan and limited copying and communication in accordance with the *Copyright Act 1968*.



Kim Jye Lee Chang

August 2013

Table of Contents

Declaration	ii
List of Tables	vii
List of Figures	ix
Abbreviations	xii
Publications	xiv
Statement of Co-Authorship	xv
Abstract	xviii
Acknowledgements	xx
Chapter 1 Introduction: Microalgae - A renewable source of biofuels, omega-3 oils and other co-products	1
Background	1
Cultivation of microalgae	2
Open versus closed cultivation systems.....	2
Potential of microalgal oil for transport fuels	5
Benefits of algal derived biofuel	9
Biodiesel potential of algae	11
Strain selection for production of biodiesel	14
Potential of microalgae for omega-3 oils and co-products	15
Rationale for this study	17
Objectives of this study.....	18
Chapter 2 Biodiscovery of new Australian thraustochytrids for production of biodiesel and long-chain omega-3 oils	20
Abstract	20
Introduction	21
Materials and methods	23
Sampling location, isolation and growth of culture organisms.....	23
Preparation and analysis of fatty acid methyl esters (FAME) and sterols	26
Extraction, amplification and sequencing of 18S ribosomal RNA genes	27
Pigment extraction and analysis.....	28
Statistical analysis.....	29
Results	30

Biodiscovery.....	30
Fatty acid and sterol profiles.....	30
18S rRNA gene sequencing.....	38
Pigment profiles.....	40
Discussion	41
Sampling location and isolation.....	41
Groupings based on 18S rRNA gene sequence, fatty acid and sterol profiles	43
Biodiversity.....	46
Omega-3 oil production.....	48
Biodiesel production.....	49
Sterols	51
Pigments.....	51
Chapter 3 Odd-chain polyunsaturated fatty acids in thraustochytrids	53
Abstract.....	53
Introduction	54
Materials and methods	56
Culture maintenance	56
Lipid extraction.....	57
Preparation of 4,4-dimethyloxazoline (DMOX) derivatives.....	57
GC and GC-MS analysis.....	58
Results	59
Discussion	64
Chapter 4 Comparison of thraustochytrids <i>Aurantiochytrium</i> sp.,	
<i>Schizochytrium</i> sp., <i>Thraustochytrium</i> sp. and <i>Ulkenia</i> sp. for production of	
biodiesel, long-chain omega-3 oils and exopolysaccharides.....	70
Abstract.....	70
Introduction	71
Materials and methods	74
Microorganisms and culture condition	74
Biomass harvest and analysis of fatty acid methyl esters (FAME).....	76
Harvesting and characterisation of polysaccharides.....	77
Results and Discussion	78
Effect of glucose on biomass yield by different thraustochytrid strains.....	78
Correlation of growth and pH change	86
Biomass and fatty acid profiles	87

Yield and crude chemical characterization of exopolysaccharides.....	94
Conclusion.....	98
Chapter 5 High cell density cultivation of a novel <i>Aurantiochytrium</i> sp. strain TC 20 in a fed-batch system using glycerol to produce feedstock for biodiesel and omega-3 oils	99
Introduction	100
Materials and methods	105
Microorganism and culture conditions	105
Biomass harvest and analysis of fatty acid methyl esters (FAME).....	108
Results	110
Effect of initial glucose levels (40 g/L and 100 g/L) on cell growth and lipid yield	110
Evaluation of glycerol as carbon source.....	113
Glycerol and additional nutrients in fed-batch cultivation.....	116
Discussion	119
Effect of initial glucose levels on biomass and fatty acid production.....	119
Comparison of glucose and glycerol as carbon source	121
Glycerol and additional nutrients in fed-batch cultivation.....	122
pH regulation of <i>Aurantiochytrium</i> sp. TC 20 in fermentation.....	123
Biodiesel and long-chain omega-3 fatty acid production	124
Chapter 6 Life-cycle assessment: Heterotrophic cultivation of thraustochytrids for biodiesel production.....	128
Abstract.....	128
Introduction	129
Materials and methods	133
LCA software	133
Life cycle assessment.....	133
Heterotrophic cultivation	136
Processing	138
Combustion emissions	140
Results and Discussion	142
Total greenhouse gas impact and energy intensity.....	142
Conclusion.....	147
Chapter 7 Conclusions and future perspectives	148

Introduction	148
Biodiscovery	149
Odd-chain polyunsaturated fatty acids	149
One litre batch culture trials	151
Optimisation in two litre fed-batch fermentation trials in bioreactors.....	151
Life-cycle assessment	152
Fatty acid profiles.....	153
Biofuels, omega-3 oils and co-products	155
Implications for further research	157
Bibliography	158

List of Tables

Table 1-1 Advantages and limitations of open ponds and photobioreactors (adapted from Brennan and Owende, 2010).....	4
Table 1-2 Major investments during 2008 - 2009 in biotech companies to develop renewable liquid fuels from microalgae in USA (summarised from Mascarelli, 2009b)	7
Table 1-3 Oil content of selected microalgae (data from Chisti, 2007).....	12
Table 2-1 Pigment composition and content ($\mu\text{g/L}$) of selected strains from each taxonomic group (based on fatty acid and sterol profiles).....	40
Table 2-2 Summary of chemotaxonomic characteristics and 18s rRNA sequence data of thraustochytrid strains examined in the present study.....	44
Table 3-1 Fatty acid (FA) composition (%) of two thraustochytrid strains (TC 01 and TC 04) isolated from the south east coast of Tasmania (n = 2).....	60
Table 3-2 Characteristic and diagnostic fragment ions from DMOX derivatives of selected fatty acids from thraustochytrids TC 01 and TC 04.....	62
Table 4-1 List of thraustochytrid strains and their chemotaxonomic and molecular grouping, based on Lee Chang et al. (2012)	74
Table 4-2 Fatty acid composition (as percentage of total fatty acids) for eight thraustochytrid strains grown in 4 % w/v glucose in shake flask culture at day 9	90
Table 4-3 EPS yield at day 7 for eight thraustochytrid strains grown in shake flask cultures with 2 % w/v glucose	95
Table 5-1 Fatty acid profiles (as either % of total FA or mg/g biomass) of <i>Aurantiochytrium</i> sp. strain TC 20 (CS number: CS-980, GenBank accession numbers: JN675267) at 69 h of fermentation in different fed-batch treatments of experiment 2: (i) glucose (Glu); (ii) glycerol (Gly) and (iii) glycerol and nutrients (Gly/ Nutr)	125
Table 5-2 Lipid class composition (as % of total lipids) of <i>Aurantiochytrium</i> sp. strain TC 20 (Gly/ Nutr) at 69 h of fermentation with glycerol as the carbon source. Composition determined using thin layer chromatography – flame ionisation detection.....	126

Table 6-1 Bioreactor inputs (operating conditions) and outputs (microalgal culture) per 0.20 ML reactor batch, used for the LCA (glycerol and outputs scaled up from data in Lee Chang et al., 2013).	137
Table 6-2 Process inputs (operational parameters) and outputs (products) per 0.20 ML reactor batch, used for the LCA.....	139
Table 6-3 Different greenhouse gases and their global warming potentials (GWP) relative to carbon dioxide (with a 100 year time horizon).....	141
Table 7-1 Fatty acid profiles (as percentage of total fatty acids) of <i>Aurantiochytrium</i> sp. strain TC20 (Group F) from three different experiments – (i) Initial screening, (ii) Flask optimization, and (iii) Scale up in bioreactors	154

List of Figures

- Figure 2-1 Chemotaxonomic grouping of 36 strains of temperate (*) and tropical (^) new Australian thraustochytrids based on fatty acid and sterol profiles formed by using the complete linkage function to compare Bray-Curtis similarity matrices (Spearman correlation, 9999 permutations) in Primer 6 (PRIMER-E Ltd, Plymouth, UK).....31
- Figure 2-2 Phylogeny of thraustochytrid partial 18S rRNA gene sequences from DNA extracts of the 36 new Australian strains (* temperate strains, ^ tropical strains) and reference sequences from the database. Parentheses indicate sequences matched to chemotaxonomic groups. Bootstrap values > 75 % are shown at the nodes. Two strains which were chemotaxonomically grouped as E, clustered more closely with other groups. These were TC 08 (clustered with groups G and H) and TC 07 (clustered with group C). Group G strains were clustered with group H strains. Most strains in groups G and H were from tropical north Queensland, except TC 06 and TC 0839
- Figure 3-1 Mass spectra of DMOX derivatives of the two major C₂₁ PUFA: (A) 21:5 ω 5 (4,7,10,13,16) and (B) 21:4 ω 7 (5,8,11,14). Ions used to identify double bond positions at Δ 4 and Δ 5 are 152 and 153, respectively. Other double bond positions are located where a gap of 12 amu is flanked by two intense fragment ions of 40 amu difference63
- Figure 3-2 Gas chromatogram of the total FA (as FAME) of thraustochytrid TC01, showing unusual odd-chain polyunsaturated fatty acids65
- Figure 3-3 Proposed OC-FA biosynthesis pathway in thraustochytrids TC 01 and TC 04. Top row indicates number of double bonds; vertical arrows indicate elongase (e). Excluding the initial Δ 9, Δ 10 or Δ 11 desaturation, upward angled arrows indicate methylene interrupted desaturation towards the omega (ω , methyl) side of an existing double bond (as occurs in organisms which can produce PUFA de novo, including most photosynthetic and some heterotrophic organisms), downward angled arrows indicate methylene interrupted desaturation towards the delta (Δ , carboxyl) side of an existing double bond, FA in brackets were not identified in our samples, but indicate likely pathway intermediates.....68

Figure 4-1 <i>Aurantiochytrium</i> sp. strains TC 09 (Group E), 18 (Group H), 20 (Group F), and 30 (Group G) in 2 % w/v and 4 % w/v glucose. (a) biomass yield g/L, (b) glucose concentration (g/L), and (c) pH change in culture media.....	79
Figure 4-2 <i>Schizochytrium</i> sp. strain TC 02 (Group A) and <i>Ulkenia</i> sp. strain TC 10 (Group D) in 2 % w/v and 4 % w/v glucose. (a) biomass yield g/L, (b) glucose concentration (g/L), and (c) pH change in culture media	83
Figure 4-3 <i>Thraustochytrium</i> sp. strains TC 04 (Group B) and TC 33 (Group C) in 2 % w/v and 4 % w/v glucose. (a) biomass yield g/L, (b) glucose concentration (g/L), and (c) the pH change in culture media	84
Figure 4-4 The total content of (a) fatty acid methyl esters, FAME mg/g, and (b) biomass g/L, of eight different thraustochytrid strains in 2 % w/v and 4 % w/v glucose	89
Figure 4-5 The percentage of fatty acid methyl esters, FAME (a) docosahexaenoic acid, DHA, and (b) palmitic acid, 16:0, of eight different thraustochytrid strains in 2 % w/v and 4 % w/v glucose.....	93
Figure 4-6 Crude chemical compositions of purified EPS produced by thraustochytrid strains grown in liquid shake flask cultures. Abundances of neutral sugars, uronic acids and protein are reported as percentage of total EPS	97
Figure 5-1 Two litres stirred tank bioreactors connected to a Biostat B (Sartorius Stedim, Australia) control system.....	106
Figure 5-2 Fed-batch cultivation of <i>Aurantiochytrium</i> sp. strain TC 20 in a 2 L bioreactor with 1.6 L of medium. The initial medium contained 40 g/L glucose. A dose of glucose (660 g/L) was fed into the bioreactor at 29 h. Data given are the mean of two replicate bioreactors under the same conditions. Abbreviations: Glu, glucose; DCW, dry cell weight.....	111
Figure 5-3 Fed-batch cultivation of <i>Aurantiochytrium</i> sp. strain TC 20 in a 2 L bioreactor with 1.6 L of medium. The initial medium contained 100 g/L glucose. Seed culture was inoculated at 5% (v/v). A dose of glucose (660 g/L) was fed into the bioreactor at 45.6 h. Data given are the mean of two replicate bioreactors under the same conditions. Abbreviations: Glu, glucose; DCW, dry cell weight.....	112
Figure 5-4 Fed-batch cultivation of <i>Aurantiochytrium</i> sp. strain TC 20 in a 2 L bioreactor with 1.6 L of medium. The initial medium contained 40 g/L glucose (Glu) or glycerol (Gly). An additional 400 mL 660 g/L of glucose (Glu) or	

glycerol (Gly) was fed into the bioreactor at 25.4 h. Abbreviations: DCW, dry cell weight.....	114
Figure 5-5 Effect of addition of nutrients (Gly/ Nutr) in a fed-batch cultivation of <i>Aurantiochytrium</i> sp. strain TC 20 in a 2 L bioreactor with 1.6 L of medium. The initial medium contained 40 g/L glycerol. At 25.4 h, an additional feed of 400 mL 660 g/L glycerol + 100 mL of concentrated nutrients (yeast extract, trace metal mix, vitamin mix, MgSO ₄ .7H ₂ O and MSG) was added at 18 mL/h (total volume of fermentor = 2 L). Abbreviation: DCW, dry cell weight.....	115
Figure 5-6 Yield of (a) saturated fatty acid, 16:0, and (b) docosahexaenoic acid, DHA, in different treatments of the <i>Aurantiochytrium</i> sp. strain TC 20. All 3 treatment media were initially with 40 g/L of glucose (Glu) or glycerol (Gly). An additional 400 mL of 660 g/L glucose or glycerol was fed at 25.4 h. Bioreactor Gly/ Nutr had an additional 100 mL of concentrated nutrients (yeast extract, trace metal mix, vitamin, MgSO ₄ .7H ₂ O and MSG) added at 25.4 h.....	117
Figure 5-7 Partial gas chromatogram of the total FA (as FAME) of the thraustochytrid <i>Aurantiochytrium</i> sp. strain TC 20 at 69 h of fermentation with glycerol as the carbon source (Gly/ Nutr). The shoulder on the front of DHA is DPA-6. Abbreviations: AA, 20:4ω6; EPA, 20:5ω3; DPA-6, 22:5ω6; DHA, 22:6ω3	118
Figure 6-1 Heterotrophic microalgae production process diagram. The upgrade step could be production of biodiesel via either hydroprocessing to produce a hydrocarbon based fuel, or transesterification to produce a fatty acid methyl ester based fuel. The residual biomass could be used elsewhere for example for stock or aquaculture feed or used on-site in an anaerobic digester to produce energy to be fed back into the system.....	135
Figure 6-2 LCA of biodiesel derived from heterotrophic microalgae production followed by extraction and transesterification and energy yields resulting from oil-free biomass back to anaerobic digestion.....	144

Abbreviations

The following abbreviations have been used in this thesis:

15:0	pentadecanoic acid
16:0	palmitic acid
17:0	heptadecanoic acid
18S rRNA	18S ribosomal RNA gene
AA	arachidonic acid (20:4 ω 6)
ANACC	Australian National Algae Culture Collection
AQIS	Australian Quarantine and Inspection Service
BSTFA	N,O-Bis(trimethylsilyl)trifluoroacetamide
DCW	Dry cell weight
DHA	docosahexaenoic acid (22:6 ω 3)
DMOX	4,4-dimethyloxazoline
DPA-3	docosapentaenoic acid (ω 3)
DPA-6	docosapentaenoic acid (ω 6)
EPA	eicosapentaenoic acid (20:5 ω 3)
EPS	exopolysaccharides
ERoEI	energy returned on energy invested
FAME	fatty acid methyl ester
FAS	fatty acid synthase
FID	flame ionization detector
GC	gas chromatography
GC-MS	gas chromatography- mass spectrometry
Glu	glucose
Gly	glycerol
HC	hydrocarbon
HCl	hydrochloric acid
HPLC	high performance liquid chromatography
HRD	hydroprocessed renewable biodiesel
LCA	Life-cycle assessment

LC-PUFA	long chain ($\geq C_{20}$) polyunsaturated fatty acid
MeOH	methanol
MUFA	monounsaturated fatty acid/s
Nutr	nutrients
OC-FA	odd-chain fatty acids
PCR	polymerase chain reactions
PKS	polyketide synthase
PUFA	polyunsaturated fatty acid/s
SD	standard deviation
SFA	saturated fatty acid
SPI	septum-programmable injector
TAG	triacylglycerols
TC	thraustochytrids collection
TFA	total fatty acid/s
Tr	trace
X:Y ω Z	This was adopted for the naming of fatty acids, where X refers to the number of carbon atoms in the molecule, Y refers to the number of double bonds in the molecule, and Z indicates the carbon position of the first double bond from the terminal methyl end (CH ₃) of the molecule. The latter is generally referred to as omega Z (e.g. ωZ) or n-Z (e.g. n minus Z).

Publications

The following peer-reviewed publications have been either wholly or partially derived from work associated with this thesis:

Kim Jye Lee Chang, Maged P. Mansour, Graeme A. Dunstan, Susan I. Blackburn, Anthony Koutoulis and Peter D. Nichols, **Novel odd-chain polyunsaturated fatty acids in thraustochytrids**, *Phytochemistry* 72 (2011) 1460–1465

Kim Jye Lee Chang, Graeme A. Dunstan, Guy C.J. Abell, Lesley A. Clementson, Susan I. Blackburn, Peter D. Nichols and Anthony Koutoulis, **Biodiscovery of new Australian thraustochytrids for production of biodiesel and long-chain omega-3 oils**, *Applied Microbiology and Biotechnology* (2012) 93:2215–2231

Kim Jye Lee Chang, Geoff Dumsday, Peter D. Nichols, Graeme A. Dunstan, Susan I. Blackburn and Anthony Koutoulis, **High cell density cultivation of a novel *Aurantiochytrium* sp. strain TC 20 in a fed-batch system using glycerol to produce feedstock for biodiesel and omega-3 oils**, *Applied Microbiology and Biotechnology* (2013) 97:6907–6918

Kim Jye Lee Chang, Carol Mancuso Nichols, Susan I. Blackburn, Graeme A. Dunstan, Anthony Koutoulis and Peter D. Nichols, **Comparison of thraustochytrids *Aurantiochytrium* sp., *Schizochytrium* sp., *Thraustochytrium* sp. and *Ulkenia* sp. for production of biodiesel, long-chain omega-3 oils and exopolysaccharide**, manuscript submitted to *Marine Biotechnology* (18 Dec 2012).

Kim Jye Lee Chang, Lucas Rye, Tim Grant, Graeme A. Dunstan, Susan I. Blackburn, Peter D. Nichols and Anthony Koutoulis, **Life-cycle assessment: Heterotrophic cultivation of thraustochytrids for biodiesel production**, manuscript in preparation.

Statement of Co-Authorship

The following people and institutions contributed to the publication of work undertaken as part of this thesis:

Candidate: Kim Jye Lee Chang, School of Plant Science

Author 1: Anthony Koutoulis, UTAS School of Plant Science

Author 2: Susan I. Blackburn, CSIRO Marine and Atmospheric Research

Author 3: Peter D. Nichols, CSIRO Marine and Atmospheric Research

Author 4: Graeme A. Dunstan, CSIRO Marine and Atmospheric Research

Author 5: Maged P. Mansour, CSIRO Marine and Atmospheric Research

Author 6: Guy C.J. Abell, CSIRO Marine and Atmospheric Research

Author 7: Lesley A. Clementson, CSIRO Marine and Atmospheric Research

Author 8: Carol Mancuso Nichols, CSIRO Materials Science and Engineering

Author 9: Geoff Dumsday, CSIRO Materials Science and Engineering

Author 10: Lucas Rye, CSIRO Marine and Atmospheric Research

Author 11: Tim Grant, Life Cycle Strategies

Author details and their roles:

Paper 1, Novel odd-chain polyunsaturated fatty acids in thraustochytrids:

Located in chapter 3. Candidate was the primary author and author 3, author 4 and author 5 contributed to the lipid identification and laboratory assistance.

Authors 1,2, 3, and 4 assisted with refinement and presentation.

Paper 2, Biodiscovery of new Australian thraustochytrids for production of biodiesel and long-chain omega-3 oils: *Located in chapter 2. Candidate was the primary author, author 3 and author 4 contributed to the lipid identification, author 6 assisted with DNA extraction and sequencing software, and author 7 contributed to the pigment isolation and analysis. Authors 1, 2, 3, and 4 assisted with refinement and presentation.*

Paper 3, Comparison of thraustochytrids *Aurantiochytrium* sp., *Schizochytrium* sp., *Thraustochytrium* sp. and *Ulkenia* sp. for production of biodiesel, long-chain omega-3 oils and exopolysaccharides: *Located in chapter 4. Candidate was the primary author, author 3 and author 4 contributed to the lipid identification, and with author 8 contributed to the EPS isolation and characterisation. Authors 1, 2, 3, and 4 assisted with refinement and presentation.*


Paper 4, High cell density cultivation of a novel *Aurantiochytrium* sp. strain TC 20 in a fed-batch system using glycerol to produce feedstock for biodiesel and omega-3 oils: *Located in chapter 5. Candidate was the primary author, author 3 and author 4 contributed to the lipid identification, and with author 9 contributed to the experimental design and Bioreactor operation. Authors 1, 2, 3, and 4 assisted with refinement and presentation.*

Paper 5, Life-cycle assessment: Heterotrophic cultivation of thraustochytrids for biodiesel production: *Located in chapter 6. Candidate was the primary author, author 10 and author 11 contributed to the idea, LCA software operation and development. Authors 1, 2, 3, and 4 assisted with refinement and presentation.*

We the undersigned agree with the above stated "proportion of work undertaken" for each of the above published (or submitted) peer-reviewed manuscripts contributing to this thesis:

Signed: 

Anthony Koutoulis
Supervisor
School of Plant Science
University of Tasmania




René Vaillancourt
Head of School
School of Plant Science
University of Tasmania



Susan I. Blackburn
Supervisor
Marine and Atmospheric Research
CSIRO



Peter D. Nichols
Supervisor
Marine and Atmospheric Research
CSIRO



Graeme A. Dunstan
Supervisor
Marine and Atmospheric Research
CSIRO

Date: 7 August 2013

Abstract

The potential of biofuel production from microalgae is of intense interest globally owing to growing concern with rising crude oil prices and future availability. In addition to producing lipids for potential biofuel application, thraustochytrids are capable of forming other high-value bioproducts, such as proteins, enzymes, omega-3 polyunsaturated fatty acids (PUFA), carotenoid pigments and exopolysaccharides (EPS). The co-production of high-value bioproducts during biofuel production is desirable when it adds greater value to the production process and improves process economics.

Thirty-six new thraustochytrids have been isolated from the southeast coast of Tasmania and far north Queensland. They were separated into eight chemotaxonomic groups (A – H) based on fatty acid and sterol composition, with the groups clustered closely with four different genera (*Aurantiochytrium*, *Schizochytrium*, *Thraustochytrium* and *Ulkenia*) based on 18S rDNA molecular identification. In an initial screening study, some strains produced > 60 % docosahexaenoic acid (DHA) under unoptimized culture conditions. *Aurantiochytrium* sp. strains (groups G and H) contained 15:0 (pentadecanoic acid) at between 20 – 30 % of the total fatty acids (TFA) and 16:0 (palmitic acid) in the range of 7 – 15 % TFA, suggesting these strains could be potential candidates for biodiesel production. β,β -Carotene, canthaxanthin and astaxanthin were identified in pigmented strains. Part of the process to scale up is to select the best performing strain based on growth and biochemical characteristics. In the subsequent trials, eight thraustochytrid strains from the different chemotaxonomic groups (A – H) were compared in 1 L scale baffled shake flasks for the synthesis of EPS, in addition to biomass yield and fatty acid profiles. The crude chemical characterization of the EPS, which were released into the culture media by these strains, was performed as an initial step in

determining the potential for biotechnological application of these biomaterials. *Aurantiochytrium* sp. strain TC 20 had the highest biomass production (18.5 g/L) and oil yield (7.5 g/L) after 9 days of growth in 4 % w/v glucose basal media at 20 °C, with 0.18 g/L EPS extracted from the supernatant. The maximum yield of EPS was observed in *Schizochytrium* sp. strain TC 02 (0.3 g/L). High biomass producing strains that also had high lipid and high EPS yield may be better candidates for commercial production of biofuels and other bioproducts. The next phase was to optimize biomass in 2 L bioreactors. The growth of *Aurantiochytrium* sp. TC 20 was also investigated using glycerol as a carbon source. Glycerol is becoming increasingly available, because it is a by-product of biofuel production from vegetable oil and animal fats. Fortification of the feed with additional nutrients improved the biomass yield from 56 g/L (34 % total fatty acids) to 71 g/L (52 % total fatty acids, cell dry weight) at 69 h.

A life-cycle assessment, from the upstream biomass production to the direct emission of biodiesel combustion, was applied to assess the energy balance and the potential environmental impacts of this heterotrophic microalgal-derived biodiesel. The scenario analysis of a virtual production facility, modeled on experimental yield data, demonstrated that cultivation of heterotrophic microalgae for the production of biodiesel is comparable in terms of greenhouse gas emissions and energy usage to production of petroleum diesel. The LCA identified that improvements in cultivation conditions, in particular the bioreactor energy inputs and microalgae yield, will be critical in developing a sustainable production system. This study demonstrates the potential of heterotrophic cultivation of newly isolated endemic thraustochytrids to provide Australia's transportation fleet with a secure, environmentally sustainable alternative fuel feedstock, and co-production of high value bioproducts that can provide additional revenue to benefit the economics of biofuel production.

Acknowledgements

I would like to acknowledge the following people who helped complete my thesis. Foremost should be my supervisors, Anthony Koutoulis, Susan Blackburn, Peter Nichols and Graeme Dunstan, all of whose invaluable knowledge and guidance made this journey stimulating and enjoyable. I am blessed to work with them and they have shown endless patience and made important comments during the thesis writing. I am grateful to Geoff Dumsday and Carol Nichols for providing invaluable input during the fermentation study.

I would like to thank all those in CSIRO who have so kindly offered their time, assistance on various matters, in particular Malcolm Brown, Lesley Clementson, Guy Abell, Ian Jameson, Peter Mansour, Lucas Rye, Dave Batten, Tim Grant, Dion Frampton, Cathy Johnston, Sharon Appleyard, Elisabeth Albinsson, Tim Fountain, Rob Gurney, Mina Brock, Danny Holdsworth and Birgit Unterweger.

In addition, I am indebted to the Australian government for the Australian Postgraduate Award (APA) and the CSIRO Office of the Chief Executive (OCE) top-up scholarship through the CSIRO Energy Transformed Flagship for funding my PhD study.

My family and friends have been equally supportive and any accomplishments of mine are due in no small part to their company. They include Wang Heong Lee, Moi Yeng Chiang, Kim Par Lee Chang, Rachel Yong, Don Soh, Shiok Cheng Lee, Shwuyi Patulski, Kurt Patulski, David Keys, Shelley Keys, Ming Teck Lim, Bob Lavis, Michael Carnes and Eric Hsu.

Last but not least, Sia Wee Lim, my girlfriend, for replenishing my soul.