Key Points:

- We developed a new method to estimate vertical land motion and regional altimeter (ALT) systematic errors, tested in the Baltic Sea region
- We detected localized motion at several gauges in the Baltic region that cannot be inferred from interpolated global positioning system or geophysical model velocities
- We captured significant ALT errors and improved the agreement between absolute sea-level from tide gauge and altimetry records

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

M.-H. Rezvani,
mohammadhadi.rezvani@utas.edu.au

Citation:

Rezvani, M.-H., Watson, C. S., & King, M. A. (2021). Estimating vertical land motion and residual altimeter systematic errors using a Kalman-based approach. *Journal of Geophysical Research: Oceans*, 126, e2020JC017106. <https://doi.org/10.1029/2020JC017106>

Received 19 DEC 2020

Accepted 21 MAY 2021

Estimating Vertical Land Motion and Residual Altimeter Systematic Errors Using a Kalman-Based Approach

Mohammad-Hadi Rezvani¹ , Christopher S. Watson¹ , and Matt A. King¹

¹School of Geography, Planning, and Spatial Sciences, University of Tasmania, Hobart, TAS, Australia

Abstract Vertical land motion (VLM) is the connection between absolute sea-level (ASL) from a satellite altimeter (ALT) and relative sea-level from a tide gauge (TG). VLM is often sparsely observed yet is required for understanding sea-level rise. Many studies have sought to exploit ALT and TG data to infer VLM, yet regionally correlated systematic errors in altimetry have not been considered. We have developed a Kalman filtering and smoothing framework to simultaneously estimate location-specific VLM and residual mission-specific systematic errors in a geocentric reference frame. We used ALT minus TG, ALT crossovers and global positioning system (GPS) bedrock height observations in a multi-stage solution approach that gradually separated time-variable parameter estimates in an ill-posed problem. We evaluated the performance of the method using the Jason-series along-track data in the Baltic Sea, where glacial isostatic adjustment is the dominant driver of VLM. We estimated local VLM variability at TGs of up to ~4.5 mm/yr which is not evident in spatially interpolated GPS velocities. The estimated regional altimeter errors are significant and within the range of ~±0.5–2.5 mm/yr. Our approach improves agreement between ASL estimates from ALT and TG records, provides a ~20% decrease in root mean squared error of latitudinal ASL variability at TGs, and a reduction of the ASL rate from altimetry by ~0.3 mm/yr across the region. This method advances the ALT-TG approach to determining VLM at TG locations and systematic errors of altimetry, which is broadly applicable to other regional- and global-scale studies.

Plain Language Summary Changes in sea-level can be measured relative to the land by tide gauges, or relative to the center of mass of the Earth by satellite altimeters. These two different types of sea-level measurements will differ if the land is moving (vertical land motion (VLM)), or if there are errors in either system. Here we compared tide gauge and altimeter data to estimate how much the land is moving - at the same time, we estimated small but significant errors in the altimetry system at a regional scale. We tested our new method over the Baltic Sea and showed this revealed local motion of some tide gauges that could not be previously detected using other approaches, such as interpolation of VLM determined from nearby global positioning system sites. Our approach helps refine sea-level change measurements which are critical for informed management of coastal regions.

1. Introduction

Vertical land motion (VLM) of the Earth's crust occurs over diverse spatial and temporal scales, driven by surface mass-loading, anthropogenic influences, and tectonic processes (e.g., Pfeffer et al., 2017). VLM provides the key connection between relative sea-level (RSL) derived from a tide gauge (TG) and absolute sea-level (ASL) derived from a satellite altimeter (ALT, e.g., Church & White, 2011). ASL data is mainly comprised of ALT-inferred time series with a near-global coverage, while RSL is typically observed along well-developed coasts (Neumann et al., 2015). In the areas with sparse long-term TG records, improved understanding of VLM is critical to help plan better sea-level rise adaptation pathways at local and regional scales (Cazenave et al., 2014; Church et al., 2013; Woodworth et al., 2011).

VLM at TG locations is obtained from geodetic observations or, in many cases, inferred from models of glacial isostatic adjustment (GIA). VLM is directly observable using space-borne techniques such as GNSS (of which global positioning system (GPS) is the most frequently used, for example, Hammond et al., 2016; Santamaría-Gómez et al., 2011), and TG VLMs are often interpolated from GPS velocities (e.g., Bouin & Wöppelmann, 2010). In the absence of nearby GPS, GIA rates are alternatively applied (e.g., Hay et al., 2015; Spada & Galassi, 2012), with an often-untested assumption that GIA is the dominant contributor to VLM.

Inferring VLM at TGs using GPS or GIA is limited. The timespan of GPS data is mostly less than a decade, and an extrapolated rate is inadequate in many locations (e.g., Featherstone et al., 2015). With a spatially sparse GPS network, there is an inevitable assumption that the TGs are not influenced by localized dynamics (e.g., Wöppelmann et al., 2009; Woodworth et al., 2017). Comparison of GPS-derived VLM with GIA signals often highlights regional discrepancies due to the presence of non-GIA processes (e.g., King et al., 2012).

Given these limitations, numerous investigations have combined ASL from multi-mission ALT and RSL from multiple TGs to estimate VLM along coastlines (e.g., early studies from Cazenave et al., 1999 and Nerem & Mitchum, 2002). Although this approach could estimate TG VLM since the early 1990s, it is inevitably affected by residual mission-specific systematic errors (e.g., Beckley et al., 2017), often referred to as bias drift (e.g., C. S. Watson et al., 2015). This may be expressed at global or regional scales (Fu & Haines, 2013) and reflects an accumulation of altimetry errors (often within mission specifications) in satellite orbits, and observed ranges, as well as mismodeled geophysical and environmental processes. The inadequacy of geophysical corrections remains a significant source of errors in the coastal zones (Benveniste et al., 2020). Geographically correlated discrepancies between various orbit solutions, reaching $\sim 2\text{--}4$ mm/yr over the operational span of a typical mission, underscore the inherent challenge of estimating sea-level trends over regional scales (e.g., Couhert et al., 2015; Fu & Haines, 2013). Such differences are greater than the VLM in many locations. Averaged globally, bias drifts for some altimeter data sets can be non-negligible (e.g., C. S. Watson et al., 2015).

Estimating VLM from combined ALT and TG data typically involves two basic assumptions (e.g., Kleinherenbrink et al., 2018; Kuo et al., 2004; Ostanciaux et al., 2012; Pfeffer & Allemand, 2016; P. J. Watson, 2019). First, VLM is parameterized using a linear model following the prevailing approach from the geodetic community. Second any regional or local expression of the altimeter-specific bias drift is considered negligible; an exception is the work of Santamaría-Gómez et al. (2014), who used double differences between pairs of nearby TGs and ALT CPs to attenuate this effect and enable the computation of relative VLM at TGs.

Here we present a new methodology centered on a space-time Kalman filter and smoother to simultaneously derive time-variable estimates of both location-specific VLM and mission-specific bias drift making the use of ALT minus TG, ALT tandem/dual crossover and GPS bedrock height measurements. We develop a flexible framework with an ability to constrain parameters over space and time, with appropriate treatment of the time-correlated noise structure and spatiotemporal covariances in our observations. Owing to the close correlation between unknowns, we gradually separate parameters within a multi-stage solution approach. We assess the performance of the method using along-track records of the reference missions spanning from September 1992 to March 2020 in the Baltic Sea, surrounded by the coasts with a dense network of TG and GPS sites.

2. Methodology

2.1. Overview

Figure 1a shows a schematic overview of the data set and nomenclature involved in our multi-stage approach, which is outlined in Figure 1b. First, 1-Hz along-track ASL from $n_{\text{ALT}} = (1, 2, \dots, l)$ altimeters were extracted at specific offshore comparison point (CP) locations, such that $n_{\text{CP}} = (1, 2, \dots, j)$. Hourly RSL from $n_{\text{TG}} = (1, 2, \dots, i)$ TGs were then differenced against the contemporaneous ASL at CPs within a distance threshold to each TG (e.g., 150 km), hereon given the designation altimeter minus tide gauge (ATG). The ~ 9.9 -day sampling of the Jason-series missions was used as the time base defining each step of our Kalman computations.

Second, ASL series from multiple missions were differenced at tandem CPs, hereon designated the altimeter crossover as AXO. Tandem crossovers pertain to two missions in formation-flight on the same ground-track but separated in time by $\sim 1\text{--}2$ minutes. Dual crossovers, occurring between missions on different ground tracks (e.g., Jason-1 and Envisat), were not used given this study was limited to the Jason-series missions.

Figure 1. (a) Schematic overview of the problem and data set. The altimeter (ALT) observes absolute sea-level (ASL) at offshore comparison points (CPs, blue circles). Tide gauges (TGs, red triangles) observe relative sea-level (RSL) at the coasts. We formed altimeter minus tide gauge (ATG) observations by taking the difference of a TG/CP pair (e.g., $[z_q^{ATG}]_{CP_1, TG_1}^{ALT_1} = [z_q^{ALT}]_{CP_1}^{ALT_1} - [z_q^{TG}]_{TG_1}$). Also shown are nearby global positioning system (GPS) sites (green squares) and indicative covariances within each observational set (i.e., ASL, RSL and GPS height). A subset of tandem CPs includes altimeter crossovers (AXOs, light blue circles) where ASL differences were formed between missions ALT_1 and ALT_2 (e.g., $[z_q^{AXO}]_{CP_4}^{ALT_{1,2}} = [z_q^{ALT}]_{CP_4}^{ALT_2} - [z_q^{ALT}]_{CP_4}^{ALT_1}$). (b) Flowchart of our multi-stage solution approach to decorrelate components of the state vector in successive solutions resulting in simultaneous estimates of bias drift and vertical land motion (VLM) trend in the same reference frame.

Third, daily height series from $n_{GPS} = (1, 2, \dots, k)$ GPS sites were incorporated once resampled to the measured time of the nearest ASL data. Additionally, a priori estimates of state parameters such as VLM or altimeter drift were constrained as pseudo-observations (PSOs), depending on the specific stage of the multi-solution approach (Figure 1b). Spatiotemporal covariances within each observational set were also considered. Hereon, the terms systematic error and bias drift are used interchangeably.

Over a study region, these yielded n_{ATG} ASL minus RSL observations (z_q^{ATG}), n_{AXO} ASL differences at crossovers (z_q^{AXO}), n_{GPS} GPS heights (z_q^{GPS}) and n_{PSO} PSO constraints (z_q^{PSO}) to form our input measurement vector:

$$z_q = [z_q^{ATG} \ z_q^{AXO} \ z_q^{GPS} \ z_q^{PSO}]^T \quad (1)$$

Using the observational models described in the supporting information (see Text S1.1), we formed the state vector:

$$x_q = [r_q \ v_q \ s_q \ \alpha_q \ \psi_q]^T \quad (2)$$

comprised of (a) “time-variable” unknowns that included mission-specific bias drifts (r_q), location-specific VLM trends at TG and GPS sites (v_q), and temporally correlated measurement noise (ψ_q), as well as (b) “time-fixed” across-track ASL slopes (s_q). Additionally, we treated the intercepts (α_q), defined at the initial epoch of each observational series, as “time-variable” unknowns in the first solution of multi-stage approach, and as “time-fixed” quantities in the subsequent solutions (Figure 1b). Hereon, superscripts refer to observational sets (i.e., ATG, AXO, GPS and PSO) and the subscript q refers to the Kalman timestep of ~ 9.9 days.

We assumed that VLMs (over the lifespan of TG or GPS sites) and bias drifts (over the lifespan of respective missions) are constant in time during the first and second steps of the multi-stage strategy. However, we let them vary in time to capture their respective evolution in the last step.

We performed preliminary data analyses to provide (a) a priori estimates of unknowns, (b) content of time-invariant and -variant noise present in individual observational data, driven by environmental and instrumental errors, and (c) spatiotemporal correlations within observations. The noise characteristics along with the covariances reflect the underlying random field in the consecutive adjustments.

2.2. Kalman Framework and Implementation

The Kalman algorithm was implemented through a recursive process of forward filtering and backward smoothing. In the forward pass, a dynamic model predicted a priori estimates of the state vector and covariances using previous a posteriori estimates. The a priori estimates were then filtered using observations up to and including the epoch of computation. In the backward pass, the a posteriori estimates were optimally smoothed using all observations available over the computational timespan; this means that all estimates benefited from all observations in time.

In the forward filtering, following the method set out in Grewal and Andrews (2008), the measurement equation was constructed to relate the state vector $\mathbf{x}_q$ with the measurement vector $\mathbf{z}_q$ allocated to the time-step q as:

$$\mathbf{z}_q = \mathbf{H}_q \mathbf{x}_q + \boldsymbol{\varepsilon}_q \quad (3)$$

with design matrix $\mathbf{H}_q = \partial \mathbf{z}_q / \partial \mathbf{x}_q$ that was specified on an epoch-by-epoch basis. The Gaussian white noise of the observations $\boldsymbol{\varepsilon}_q = [\varepsilon_q^{\text{ATG}} \ \varepsilon_q^{\text{AXO}} \ \varepsilon_q^{\text{GPS}} \ \varepsilon_q^{\text{PSO}}]^T$ was comprised of individual noise pertaining to ASL minus RSL ($\varepsilon_q^{\text{ATG}}$), ASL differences at crossovers ($\varepsilon_q^{\text{AXO}}$), GPS heights ($\varepsilon_q^{\text{GPS}}$) and PSO constraints ($\varepsilon_q^{\text{PSO}}$), each with zero-mean. These elements constituted the measurement variance-covariance (VCV) matrix $\mathbf{R}_q$ that is simply expressed in a constant form $\mathbf{R}$ as:

$$\text{diag}(\mathbf{R}) \equiv \text{diag}(\mathbf{R}_q) = \left[\mathbf{R}(\varepsilon_q^{\text{ATG}}) \ \mathbf{R}(\varepsilon_q^{\text{AXO}}) \ \mathbf{R}(\varepsilon_q^{\text{GPS}}) \ \mathbf{R}(\varepsilon_q^{\text{PSO}}) \right]^T \quad (4)$$

where diagonal elements of $\mathbf{R}(\varepsilon_q^{\text{ATG}})$, $\mathbf{R}(\varepsilon_q^{\text{AXO}})$ and $\mathbf{R}(\varepsilon_q^{\text{GPS}})$ included white noise variances of the ATG, AXO and GPS records, respectively. A priori values of these time-invariant metrics were determined from spectral analysis of observation residuals following removal of a linear fit (see Section 3.3.2). The off-diagonal elements of these VCV matrices were filled with covariances inferred from empirical semi-variograms computed from the respective residuals, as described in Section 3.3.3. The diagonal $\mathbf{R}(\varepsilon_q^{\text{PSO}})$ was comprised of variances associated with tight or loose constraints.

In the forward prediction step, the state vectors at two consecutive timesteps q and $q-1$ were transitioned through the discrete-time linear dynamic model expressed as (e.g., Grewal & Andrews, 2008):

$$\mathbf{x}_q = \boldsymbol{\Phi} \mathbf{x}_{q-1} + \boldsymbol{w}_q \quad (5)$$

where $\boldsymbol{\Phi}$ is the diagonal transition matrix, equivalent to an identity matrix if all components of the state vector are treated as random-walk processes. As we approximated the time-dependent noise of observations ($\boldsymbol{\Psi}_q$) via a first-order autoregressive (AR1) stochastic model, the corresponding elements in this matrix were filled with the AR1 coefficients estimated from the spectral analysis of the residuals (see Section 3.3.2). The term $\boldsymbol{w}_q$ represents the Gaussian process noise sequences such that $\boldsymbol{w}_q = [\boldsymbol{w}_q^r \ \boldsymbol{w}_q^v \ \boldsymbol{w}_q^s \ \boldsymbol{w}_q^a \ \boldsymbol{w}_q^w]^T$ (noting superscripts here refer to parameter types). This process noise is vital to our approach in tuning unknowns including bias drifts ($\boldsymbol{w}_q^r$), VLM trends ($\boldsymbol{w}_q^v$), ASL slopes ($\boldsymbol{w}_q^s$), intercepts ($\boldsymbol{w}_q^a$) and time-correlated noise ($\boldsymbol{w}_q^w$), each with zero-mean, following the state VCV matrix $\mathbf{Q}_q$ such that:

$$\text{diag}(\mathbf{Q}) \equiv \text{diag}(\mathbf{Q}_q) = \left[\mathbf{Q}(\mathbf{w}_q^r) \mathbf{Q}(\mathbf{w}_q^v) \mathbf{Q}(\mathbf{w}_q^s) \mathbf{Q}(\mathbf{w}_q^a) \mathbf{Q}(\mathbf{w}_q^w) \right]^T \quad (6)$$

with the diagonal matrices accounted for process noise variances of bias drifts $\mathbf{Q}(\mathbf{w}_q^r)$, VLMs $\mathbf{Q}(\mathbf{w}_q^v)$, ASL slopes $\mathbf{Q}(\mathbf{w}_q^s)$, intercepts $\mathbf{Q}(\mathbf{w}_q^a)$, and time-correlated noise $\mathbf{Q}(\mathbf{w}_q^w)$.

The diagonal elements of $\mathbf{Q}(\mathbf{w}_q^w)$ were estimated from AR1 noise modeling of the residuals using a power spectrum approach (see Section 3.3.2). Here we assumed no spatial correlations within the unknowns, and we therefore considered all state VCV matrices as simply diagonal. However, where a priori knowledge of covariances between a specific set of unknowns is reliably available, such as between GPS- and TG-specific VLM or within mission-specific bias drifts or CP-specific ASL slopes, the off-diagonal elements could be populated.

We utilized the recursive relations of Kalman prediction and filtering in the forward solutions, and subsequently Rauch-Tung-Striebel smoothing in the backward solutions (Text S1.2). In practice, we first incorporated our observations (recorded with different sampling intervals at diverse times and locations) using the same timesteps that were adopted from averaging the measurement times of the reference-mission altimeters within each cycle. We thus spatiotemporally linked the noise structure and covariances within each observational set. Second, the predicted and filtered estimates of the states and the unknown covariances were determined from the first to the last computational instant. However, as some observations were not completely recorded over the full period of the satellite altimetry, we assumed individual components of the predicted and filtered states and covariances as constant in timesteps outside of the associated observational spans. Third, we used singular value decomposition to complete the inversion of the large-dimension VCV matrices in the underlying solutions. For the test study outlined below, the matrix to be inverted was typically $\sim 1900 \times 1900$ and $\sim 4000 \times 4000$ for the filter and smoother gains, respectively (see Text S1.2).

2.3. Solution Approach

We implemented the Kalman filter and smoother using the multi-stage strategy summarized in Figure 1b (see Figure S1 for more detail). Numerical values for process noise and constraints used in each solution are provided in Text S1.3.

2.3.1. Solution 1

Owing to the lack of any a priori knowledge, we used a zero-value initial state and large uncertainty for regional mission-specific bias drifts and selected a small process noise to capture subtle time-variation. We used a priori VLMs at GPS and TG sites as determined in our preprocessing steps (see Sections 3.2 and 3.3.1) as tight constraints. Owing to the plausible deviation between the nominal and actual VLMs, it is highly likely that when using time-fixed intercepts, this configuration would lead to a divergence between filtering and smoothing estimates of bias drift. To avoid this, we made the ATG intercepts time-variable using a relatively large process noise to enable the capture of both fixed-term intercepts and unmodeled VLM signals. For consistency, we adopted a similar treatment for GPS intercepts. To improve the smoothness of estimates, we iterated this solution with initial states and uncertainties updated from the median (for estimates of bias drift) and the initial-epoch (for time-varying intercepts) outputs from the first solution, respectively. In the iteration, we used trends remaining in the resultant ATG and GPS residuals to automatically adapt the process noise of intercepts to better capture any time variability present (see Table S1 for the automated rule set).

2.3.2. Solution 2

Solution 2 was achieved by placing tight constraints on PSOs pertaining to time-fixed linear bias drifts based on median estimates from the final iteration of Solution 1. We treated the intercepts as time-fixed parameters from this solution onwards. We further improved our estimates by an iteration, updating the initial VLM states and uncertainties based on median estimates from the first run. We commenced with a relatively tight process noise for VLM estimates; however, to capture localized variability in the iteration,

we made them adaptively looser according to the trends present in the resultant ATG and GPS residuals (see Table S1). We then flagged geodetic sites subject to substantial non-linear VLM by visual inspection, and thresholding of the adaptive process noise (see Figure S2). This distinguished geodetic sites with the linear motion from those with anomalous variability, so that tight or loose constraints for VLM trends in the subsequent solutions could be determined. The TGs flagged for loose constraints are listed in Text S1.3.2.

2.3.3. Solution 3

Solution 3 had a large uncertainty for initial bias drifts enabling them to adjust accordingly, and tightly constrained VLMs, for those sites identified in the previous solution as being suitably linear. A priori unknowns and uncertainties were taken based on median estimates from the former solutions. This stage was run twice (with tight constraints on the resultant bias drifts) to ensure convergence of all parameter estimates. Within this step, TGs subject to RSL datum issues were identified and excluded from the last solution (see discussion and Text S2.4.1 for further detail).

AXO intercepts were treated as tight constraints in all solutions to assist in differentiating between bias drift and VLM estimates, except for the iteration of Solution 3 where the estimates of other parameters were fixed appropriately. The across-track ASL slopes were tightly constrained to a priori values based on median estimates from the former solutions to derive time-fixed parameters. The correlated noise was also modeled with zero initial value and large uncertainty, while the respective process noise was adopted from AR1 modeling analysis (Section 3.3.2).

3. Data Set and Preliminary Analyses

We tested our approach with a case study in the Baltic region. To reduce the computational cost, we limited the number of CPs up to eight per TG, yielding 189 offshore CPs to form ATG series for five separate missions (TOPEX-side A, TOPEX-side B, Jason-1, OSTM/Jason-2 and Jason-3). We similarly limited the number of CPs to 62 over the catchment to provide AXO series in tandem phases (TOPEX-side B & Jason-1, Jason-1 & Jason-2 and Jason-2 & Jason-3).

3.1. Tide Gauge and Altimeter Data

3.1.1. Tide Gauge Relative Sea-Level

We used hourly RSL series from 41 TGs (see Figure 2) in local reference frames. These records commence from the TOPEX launch and have at least 70% completeness over the satellite era. Monthly averaged series were intercompared between neighboring TGs to check for possible outliers and datum errors. To suppress sea-level fluctuations caused by wind and atmospheric pressure (Hill et al., 2010), we applied the dynamic atmospheric correction (DAC) provided by CLS and distributed by AVISO (<https://www.aviso.altimetry.fr/>). We then estimated the small ocean tides present in the Baltic using the UTide software (Codiga, 2011). We included nodal modulations and considered the same constituents as those included within the FES2014b model (Carrere et al., 2016), thus remaining consistent with ocean tide corrections applied to ALT data (Section 3.1.2).

3.1.2. Altimeter Absolute Sea-Level

We extracted 1-Hz along-track ASL from TOPEX, Jason-1, OSTM/Jason-2 and Jason-3 archived in the Radar Altimeter Database System (RADS, Scharroo et al., 2013, data accessed March 11, 2020) relative to ITRF2008 (Altamimi et al., 2011), at CPs separated by 20 km (Figure 2). We later selected CPs within 150 km of each TG for ATG differences, while CPs spaced every 50 km were considered for tandem crossovers.

We selected GSFC-Std1204 orbits for TOPEX, and CNES-GDRE for other missions. We considered sea state bias models of Chambers-BM4 for TOPEX, CLS-nonparametric for Jason-1, and CLS-nonparametric-MLE3 for OSTM/Jason-2 and Jason-3 each using MLE3 ranges. The TOPEX data does not have the Cal-1 mode correction applied (Beckley et al., 2017). The ECMWF dry and radiometer-derived wet tropospheric corrections were selected. We applied smoothed dual-frequency ionospheric corrections, adopted with the MLE3 ranges for Jason-2 and Jason-3. For consistency, we extracted the AVISO-disseminated DAC for ALT CPs used in ATG (Section 3.1.1), and the MOG2D for AXO CPs. We applied the solid-Earth tide, pole tide,

Figure 2. Map of (a) our estimated vertical land motions (VLMs), and (b) VLM differences, global positioning system (GPS)-Krig at tide gauges (TGs) and Hector at GPS sites subtracted from our approach. TGs with VLM differences considered significant at 1-sigma are highlighted with green annotations.

FES2014a ocean load, and FES2014b ocean tide. We only corrected the radial body pole tide for ATG differences as the ocean pole tide is observed by both the TG and altimeter (Desai et al., 2015). We then estimated a priori ASL slopes at the CPs from the DTU15 mean sea surface, as a function of the zonal-track distances of each cycle-by-cycle satellite pass from the respective nominal ground tracks.

3.1.3. Altimeter Minus Tide Gauge

We linearly interpolated the RSL records at the TGs to the times of nearby altimetry observations, and then differenced to construct the ATG data set. To avoid dependence on altimeter flags, outliers specific to each CP were detected using two times the interquartile range (IQR) above or below the medians of ATG residuals following a mission-specific linear fit. The largely wind- and potentially ice-driven seasonality was then captured via harmonic analysis. We simultaneously estimated mission-specific intercepts and periodic

annual and semi-annual components specific to each CP. Removal of these seasonality terms reduced average variance in the ATG series by $\sim 34\%$.

To determine the optimal CPs, many techniques exist (e.g., Oelsmann et al., 2021). Within our CP selection process, we applied thresholds of a maximum missing data of 30% and a maximum root mean squared error (RMSE) of 0.5 m over the satellite era. Due to sea ice presence, especially in the northern regions, we considered a seemingly loose threshold of 70% missing data for ATGs over any one mission. We note that mission-specific series with at least 30% completeness can still assist in constraining VLM estimates at specific TGs, yet not cause problems with estimation of bias drifts.

We discarded the ATG series with outlying mission-specific trends that exceeded the IQR of trends from all candidate CPs for any given TG, and then selected the eight CPs with the lowest summation of white and correlated noise magnitudes (Section 3.3.2). If the ATG series for just one mission at a specific CP failed against a specific threshold, that CP was excluded from selection for any other mission. Two gauges at TRAV and KIEL locations failed to yield eight potential CPs (owing to the RMSE and gap thresholds), and were only associated with two valid CPs each. Our final data set includes 1580 ATG series from 189 CPs.

3.1.4. Altimeter Crossovers

ALT crossovers were formed by differencing the ASL from the interleaved missions. Outlying observations were detected using two times the IQR above or below the CP-specific, mission-specific medians. We imposed a minimum completeness percentage of 70% over all tandem phases and 60% within any individual phase. We then discarded within each interleaved phase, those series that exceeded two times the IQR of all white noise magnitudes in the same tandem phase (section 3.3.2). A 50 km separation criterion yielded 186 tandem crossovers at 62 CPs. Note that 32 of these CPs were used for ATG formations as well (Figure 1a).

3.2. Global Positioning System and Glacial Isostatic Adjustment Data

We used GPS daily heights from April 1994 to April 2018 provided by the Nevada Geodetic Laboratory (Blewitt et al., 2018, data accessed April 14, 2018) relative to the ITRF2008. We limited these to ensure a minimum of 3 years of data and more than 70% completeness to select 119 inland series for GPS-Krig interpolations (Table S2), later reduced to 80 sites for computational efficiency. The average span of these 80 series was ~ 9.5 years and the spatial distribution is shown in Figure 2. Outliers were detected using three times the IQR above or below the medians of the residuals following a linear fit.

We estimated a priori GPS VLM trends using the Hector software version 1.6 (Bos et al., 2013). Noise content was modeled using a “white plus power-law” stochastic model (Williams et al., 2004). We visually inspected the GPS heights to ascertain the instants of instrumental offsets and then fitted a model including intercept, offsets (where they existed), trend, and annual and semi-annual periodicities. GPS sites with a trend uncertainty larger than ± 2 mm/yr were then excluded. After removal of the offsets and seasonality terms, the resultant series were linearly interpolated at the times of the nearest ALT data to provide the 9.9156-day height series used in the Kalman filter. We used these resampled series as a sanity check on GPS VLM estimates from our Kalman solutions, in comparison with the Hector trends using the daily series.

For later comparison we linearly interpolated the ICE6G_D GIA rates (available on a $0.2^\circ \times 0.2^\circ$ grid, Peltier et al., 2018) to GPS and TG locations, last accessed November 18, 2018. This specific model was constructed up to degree and order 512 (~ 40 km) and used the VM5a Earth rheological model, ICE6G_D surface mass loading history and BEDMAP2 bathymetry in the Southern Ocean.

3.3. Preliminary Analyses

3.3.1. Relative Sea-Level and Vertical Land Motion Trends at Tide Gauge

In a similar approach to Hay et al. (2015), we estimated RSL trends at TGs using a Kalman filtering and smoothing approach with “white plus AR1” stochastic noise modeling. These RSL trends were later used in assessing the method implications for measuring sea-level rise (see Section 4.2). We derived the measurement noise magnitudes, time-correlated process noise, and AR1 transition coefficients from spectral analysis of the input data (Text S1.4.2). We set the process noise of $10^{-3}/\sqrt{9.9}$ mm/yr $\sqrt{s}$ and $10^{-1}/\sqrt{9.9}$ mm/ $\sqrt{s}$

(where s is the ~ 9.9 -day Kalman timestep) for secular trends and intercepts of all TGs. We supposed a relatively loose process noise for intercepts to appropriately derive linear trends in the presence of interdecadal variability due to the strong influence of North Atlantic Oscillation (NAO) mainly in the northern and central regions (e.g., Han et al., 2019; Hünicke & Zorita, 2008; Woolf et al., 2003). Spatiotemporal covariances within RSL noise were applied up to a length-scale of 250 km (Figure S8). We did not impose covariances within the state parameters (i.e., the RSL trends or intercepts) to avoid spatial smoothing in the presence of local variability.

Next, we estimated a priori TG VLMs using a Kriging interpolation of all GPS velocities within 150 km of each TG (hereon referred to as GPS-Krig). Note that $\sim 24\%$, $\sim 54\%$ and $\sim 78\%$ of TGs have GPS within 10, 30 and 50 km, respectively. For an individual TG, we setup a Universal Kriging model $z^{\text{VLM}} = Fx + \epsilon^{\text{VLM}}$ to link between the TG VLM unknown x and n_{VLM} GPS VLM observations z^{VLM} , through a $n_{\text{VLM}} \times 1$ matrix of ones F and Gaussian white noise ϵ^{VLM} . We interpolated the VLM trend at TG location using generalized least squares minimization (Chilès & Delfiner, 2012, page 178). We weighted the GPS trends near to an individual TG by the inverse of the white noise variances such that:

$$\left[\sigma^2(\epsilon^{\text{VLM}}) \right]_{\text{GPS}} = \left[\sigma^2(\epsilon^{\text{Nom}}) \right]_{\text{GPS}} + \left[\sigma^2(\epsilon^{\text{Dist}}) \right]_{\text{TG, GPS}} \quad (7)$$

to consider in quadrature the nominal VLM uncertainties at the GPS sites $\left[\sigma^2(\epsilon^{\text{Nom}}) \right]_{\text{GPS}}$, and added uncertainties $\left[\sigma^2(\epsilon^{\text{Dist}}) \right]_{\text{TG, GPS}}$ driven by the GPS and TG separation that were empirically inferred from semi-variogram analysis of the GPS VLM estimates over the region (see Figure S3 and Table S3).

As the semi-variogram of the noise ϵ^{VLM} can be practically approximated by the semi-variogram of the observations z^{VLM} in such a small random field (Montero et al., 2015, page 119), we computed spatial covariances between GPS VLM noise using an empirical semi-variogram of ICE6G_D predictions over Fennoscandia (Figure S4). We further added in quadrature the spatially interpolated Kriging of the GPS uncertainties (in the vicinity of 150 km) to the output Kriging uncertainty. This assisted to provide conservative uncertainties to the interpolated VLM, particularly where the GPS is only a few kilometers away from the TG (and thus the Kriging uncertainty alone would be small, Figure S5). The resultant TG uncertainties are shown in Figure S6 as a function of latitude against the nominal GPS uncertainties used in the GPS-Krig interpolations. Our a priori VLM estimates and their uncertainties at GPS and TG locations are listed in Tables S2 and S5. See Text S1.4.1 for more details.

3.3.2. Noise Analysis

To set up the Kalman filter and smoother settings, accurate knowledge of the temporally variant and invariant noise content at each measurement location is required. We used Lomb-Scargle power spectra of residuals (following removal of a linear fit) to estimate parameters quantifying the noise content of ATG and GPS observations, assuming “white plus AR1” as a plausible stochastic model. For noise at ALT tandem crossovers, we however assumed a white-only stochastic model, as the records are relatively short and ASL from two missions are measured with a delay of ~ 1 – 2 min, effectively sampling the same ocean surface.

In the case of the ATG series, we also assumed the correlated noise level to be common across all missions, as significantly contributed from the unmodeled residual ocean variability between the CP and TG locations. We fitted the following functional model to one-sided power spectrum $\mathcal{G}(\mathcal{F})$ of the observation residuals as a function of normalized frequencies, $0 < \mathcal{F} \leq 0.5$, (Buttkus, 2000; page 248):

$$\mathcal{G}(\mathcal{F}) = \sigma^2(\epsilon) + \frac{\sigma^2(w^\psi)}{1 - 2\mathcal{G}(\psi)\cos(2\pi\mathcal{F}) + \mathcal{G}(\psi)^2} \quad (8)$$

to estimate the stochastic noise parameters, which are the magnitudes $\sigma^2(\varepsilon)$ and $\sigma^2(w^\psi)$ of white and AR1 time-correlated noise respectively (where $\sigma^2(\varepsilon) < \sigma^2(w^\psi)$ is imposed), as well as AR1 coefficient $\rho(\psi)$ (where $0 < \rho(\psi) < 1$).

We populated white noise magnitudes of ATG $\sigma^2(\varepsilon^{\text{ATG}}) \equiv \sigma^2(\varepsilon_q^{\text{ATG}})$, AXO $\sigma^2(\varepsilon^{\text{AXO}}) \equiv \sigma^2(\varepsilon_q^{\text{AXO}})$ and GPS height $\sigma^2(\varepsilon^{\text{GPS}}) \equiv \sigma^2(\varepsilon_q^{\text{GPS}})$ in the matrix $\mathbf{R}_q$ to specify the associated noise levels. We populated the AR1 process noise magnitudes $\sigma^2(w^\psi) \equiv \sigma^2(w_q^\psi)$ in the diagonal elements of the matrix $\mathbf{Q}_q$ that are associated with time-correlated noise of the ATG and GPS series. The AR1 coefficients were populated as diagonal elements of the transition matrix $\mathbf{\Phi}$, that are associated with time-correlated ATG and GPS noise unknowns. For more details, see Text S1.4.2 and Figure S7.

3.3.3. Semi-Variogram Analysis

To quantify the similarity between nearby random variables at different spatial and temporal scales, we used observation covariances within each timestep of our Kalman filter. First, we separately derived empirical semi-variances of ASL ($\varepsilon_q^{\text{ALT}}$), RSL ($\varepsilon_q^{\text{TG}}$) and GPS height ($\varepsilon_q^{\text{GPS}}$) noise, by differencing the associated residuals (after removal of a linear fit) at a suite of paired locations. Second, we estimated parameters of a Gaussian negative-definite model in terms of partial sill, nugget effect and range (e.g., Montero et al., 2015), fitted to the resultant semi-variances at temporal slices (up to 10 days, Figure S8), to arrive at spatiotemporal modeling of semi-variograms and co-variograms for each set of observations. Third, we extracted the spatiotemporal covariances within ATG ($\varepsilon_q^{\text{ATG}}$) and AXO ($\varepsilon_q^{\text{AXO}}$) noise using relations from the propagation of random errors. We populated these covariances in the off-diagonal elements of the matrix $\mathbf{R}_q$ for measurements separated by up to 250 km. We therefore related the noise structure of each set of observations, recorded at different space and times, within each Kalman step. See Text S1.4.3 for more details.

4. Results

We applied the methodology to the Baltic region to derive VLM, bias drift, and subsequently ASL estimates. To evaluate the performance, we first compared our estimated VLM trends at TGs with those from the GPS-Krig interpolations and the GIA predictions. We then used our VLMs to attain ASL estimates from RSL at TGs and investigate the spatial coherence. We compared these with ASL from altimetry alone computed at CPs (with and without bias drift corrections), and findings from the solution if bias drifts were not considered.

Next, a suite of experiments was undertaken to examine the sensitivity of the solutions to unresolved issues in a priori VLM estimates and RSL datums at TGs. We then tested the sensitivity of the solution spatial domain. We also repeated our solutions to estimate bias drifts using a priori VLMs taken from the GIA model in comparison to those from the use of GPS-Krig VLMs. Finally, we assessed the Kalman performance and reliability of our results through residuals and a posteriori analysis.

4.1. Vertical Land Motion Estimates

Figure 2 shows our VLM trends estimated at 41 TG and 80 GPS sites, compared with those from the interpolated GPS-Krig and Hector analyses, respectively. A long-wavelength uplift peaked in the northwest of Bothnia and transited to maximum subsidence in the Danish straits. Our VLM trends at GPS sites are in reasonable agreement with Hector estimates with a weighted mean difference of +0.03 mm/yr and weighted root mean squared error (WRMSE) of 0.18 mm/yr (Table S4). We repeated this comparison using “white plus random-walk” modeling of the GPS height noise which resulted in a weighted mean difference of −0.01 mm/yr and WRMSE of 0.20 mm/yr. None of these VLM differences were statistically significant using either noise model. This is an important internal consistency check given we only used GPS data at the ~9.9-day Kalman timesteps (instead of daily sampling as used in Hector analysis). In contrast, our VLM estimates are significantly different from the interpolated GPS-Krig VLMs at several TGs (Table S5).

Figure 3. Profile of vertical land motion (VLM) differences per latitude, ours minus global positioning system (GPS)-Krig at tide gauges (TGs, blue squares) and ours minus Hector at GPS sites (orange circles). Significant differences at 1-sigma are labelled and highlighted in green. The weighted average and standard deviation (STD) of the differences are annotated. Error bars are ± 1 -sigma having been scaled by the a posteriori variance factor.

The presence of localized variation in VLM at some TGs could also be inferred from slightly higher uncertainties of our estimated trends, computed from Equation S21 after scaling by the a posteriori variance factor (Figure S11). We flagged TGs where the VLM differences are statistically significant at the 1-sigma level (shown in green labels in Figure 2). At the 1.96-sigma level, estimated VLMs of TGs at DAUG, RODV, SKAN and SKUL locations were identified to have significant differences to the GPS-Krig interpolated trends.

Figure 3 shows the VLM differences (ours minus GPS-Krig at TGs, and ours minus Hector at GPS sites) as a function of latitude across the region. While the GPS differences appear consistent (as expected, WRMSE of 0.18 mm/yr and standard deviation of 0.28 mm/yr), we could distinguish 16 TGs (green labels, Figures 2 and 3), where the VLMs are significantly different to GPS-Krig values (at 1-sigma). These anomalous differences range from -4.10 to $+4.59$ mm/yr. Excluding these, the standard deviation of the remaining 25 TGs is 0.70 mm/yr (reduced from 1.51 mm/yr). We repeated these comparisons against the ICE6G_D VLMs in Text S2.1.

4.2. Implications of Vertical Land Motion for Absolute Sea-Level at Tide Gauges

We estimated ASL at TGs by applying our VLM estimates to the RSL trends and investigated the spatial coherency of sea-level trends. The spatial dispersion in RSL (Figure 4a) and improved coherency in ASL (Figure 4b) suggest that we detected local variability in VLM throughout the region (at least at TG locations). Figure S12 map ASL derived using the GPS-Krig and the ICE6G_D VLMs, with substantial discrepancies from our solution persisting particularly in the Gulf of Riga. These alternative ASL show a greater variability, arguably inconsistent with the expected sea-level variation over the temporal and spatial scales considered here. Our ASL estimates at TGs (Figure 4b) are broadly consistent with those obtained from the SLCCI altimetry data set (Madsen et al., 2019), noting their analysis does not consider systematic errors in the altimetry.

Figure 5 shows a latitudinal profile of ASL estimates from our VLMs, GPS-Krig interpolations and the ICE6G_D predictions. For illustrative purposes, we modeled the variation in ASL by fitting a quadratic polynomial as a function of latitude to each data set. As shown, the difference between ASL derived from GIA and our VLM estimates steadily decrease with increasing latitude. The RMSE of the ASL trends about the corresponding polynomials is 0.87 mm/yr when using our VLMs, against 1.07 and 1.12 mm/yr from GPS-Krig and ICE6G_D VLMs, respectively, a reduction in ASL variability by $\sim 19\%$ and 22% in terms of RMSE.

As a quality control check, we derived ASL trends at already used CPs from altimetry alone, determined outside of our multi-stage approach (but still using our mission-specific bias drifts (Section 4.3) and relative intra and inter-mission biases (Text S2.3). We used an appropriate Kalman approach with “white plus AR1” stochastic noise modeling. Prior to implementation, we performed spectral analysis to determine the

Figure 4. Map of (a) relative sea-level (RSL) and (b) absolute sea-level (ASL) trends using our vertical land motion (VLM) estimates at tide gauges (TGs). As per previous figures, significant VLM differences at 1-sigma are highlighted with green annotations.

measurement noise, the process noise of time-correlated errors and AR1 transition coefficients (Text S1.4.2). We applied spatiotemporal covariances within the ASL noise from the semi-variogram analysis (Figure S8), up to a length-scale of 250 km. We estimated linear ASL trends and intercepts specific to the CPs with the same process noise used from the RSL analysis (Section 3.3.1). The process noise on the intercepts was relatively loose to withstand the effects of NAO-related variability on the secular rates (e.g., Han et al., 2019).

Figure 6 shows good agreement between our ASL estimates at TGs (using RSL and our VLMs, Table S5) and at ALT CPs (using ASL from the procedure above having applied our bias estimates). We then estimated ASL trends at the CPs from altimetry alone after applying our inter- and intra-mission biases, but not applying the bias drifts (Figure S22). The RMSE of these ASLs about the fitted polynomial is 0.78 mm/yr (compared to 0.72 mm/yr using the solution with our bias drifts and relative biases applied).

Figure 5. Profile of absolute sea-level (ASL) trends at tide gauges (TGs) by latitude, comparing estimates derived from applying our vertical land motion (VLM) to relative sea-level (RSL) (green circles) with those from global positioning system (GPS)-Krig (blue triangles) and ICE6G_D glacial isostatic adjustment (GIA) (pink squares) alternatives. Solid and dashed lines show a quadratic polynomial fitted to each set of ASL estimates per latitude with root mean squared error (RMSE) about this fit annotated in the legend. For clarity, latitudes of TGs where ASL is derived from GIA and GPS-Krig are shifted by $+0.015^\circ$ and -0.015° , respectively. Error bars are ± 1 -sigma scaled by the a posteriori variance factor.

4.3. Altimeter Systematic Errors

Figure 7 shows our estimates of mission-specific bias drift over the region. Note that the filtering (colored lines) is from the first run of Solution 3, and the smoothing (black lines) from its iteration (see Figure 1b). Convergence between the forward and backward solutions is evident after ~ 2.5 years. The averaged values of the smoothed drifts are larger than those found in global solutions (e.g., C. S. Watson et al., 2015); however, they are consistent with mission specifications and estimates of regional-specific orbit errors (e.g., Couhert et al., 2015).

As we imposed a relatively tight process noise, the estimated uncertainties for the smoothed bias drifts are likely optimistic. Alternatively, the filtering uncertainties, shown in Figure 7, are a factor of 3–4 higher and likely more realistic. The latter uncertainties are consistent with the global uncertainty assessment by C. S. Watson et al. (2019) as a function of mission duration.

Figure 6. Latitudinal profile of absolute sea-level (ASL) trends at tide gauges (TGs) using relative sea-level (RSL) plus our, global positioning system (GPS)-Krig and glacial isostatic adjustment (GIA) vertical land motion (VLM) trends, compared with those derived at comparison points (CPs) using altimetry alone after applying estimated bias drifts and relative biases. Solid and dashed lines show a quadratic polynomial fitted to each set of ASL estimates per latitude with root mean squared error (RMSE) annotated. Error bars are ± 1 -sigma scaled by the a posteriori variance factor.

Figure 7. Estimated bias drifts over the study region, estimated from Solution 3. Forward and backward estimates show convergence after approximately 2.5 years. Relatively tight process noise enables subtle evolution in time of this mission-specific parameter in the forward filtering (colored lines). The return smoothing pass effectively constrains the variables to derive time-fixed estimates (black lines and annotated values and uncertainties). The filtering uncertainties are also provided in brackets for comparison. All uncertainties are 1-sigma scaled by the a posteriori variance factor.

4.4. Sensitivity Analysis

We undertook “mild” and “extreme” perturbation experiments to assess the dependence of the solutions on a priori TG VLMs. We randomly selected six TGs, labeled SIMR, SKAG, RAUM, KEMI, FORS and LIEP (Figure 2), and artificially perturbed the ATG observations by spurious trends between -2.6 and $+2.3$ mm/yr in the mild, and between -1.4 to $+5.8$ mm/yr in the extreme, scenario (Table S7). A priori VLMs of these gauges were incorrect in comparison to the control (unperturbed) case, such that the arithmetic averages of perturbations were -0.13 and $+2.5$ mm/yr in the respective experiments. We ran the multi-stage solutions as per the control case. The results show that, on average, $\sim 82\%$ and $\sim 85\%$ of the perturbations were respectively retrieved in VLM estimates of perturbed gauges in the mild and extreme scenarios (see Figure S14). As four out of the six TGs had erroneous a priori VLMs greater than or equal to 2 mm/yr in each scenario, we expected some influence on bias drift estimates in Solution 1 given a slight shift in the “datum” imposed by the a priori TG VLMs (Figure 1b). We noticed a negligible change in bias drifts for the mild scenario (~ -0.06 mm/yr) and a small change for the extreme case ($\sim +0.31$ mm/yr). We also observed weighted average shifts in VLM estimates of unperturbed gauges of $+0.04$ and -0.24 mm/yr in the mild and extreme experiments, respectively (with WRMSE of 0.05 and 0.26 mm/yr, respectively). More details are given in Text S2.4.1.

We further investigated the worst-case scenario where the GPS network is poorly distributed, such that GPS-Krig would fail to provide reasonable a priori VLM at all TGs. We, in turn, derived a priori TG VLM outside our engine using a weighted least squares adjustment of the ATG data, constrained using null bias drifts. The resultant TG VLM estimates differed from the GPS-Krig values by a weighted average of $+0.39$ mm/yr and a relatively large standard deviation of 1.68 mm/yr. We ran the multi-stage approach using these new a priori VLMs and noticed a weighted average shift of $+0.37$ mm/yr in TG VLM trends and an associated negative change of -0.41 to -0.49 mm/yr in bias drift estimates, as per the control case. The resultant datum shift is compatible with our findings enforced with zero-drift constraints (e.g., Figure S20). This test highlights the importance of a priori VLM that is independent of altimetry systematic errors and is reasonably accurate at most TGs relative to an absolute reference frame. We return to this dependence in the discussion.

To evaluate the potential effect of RSL datum errors, we perturbed three randomly selected TGs (FORS, SIMR and SKAG, Figure 2), with artificial offsets of $+30$, $+50$ and $+10$ mm placed at the midpoint and

extending to the endpoint time of Jason-1, OSTM/Jason-2 and Jason-3 missions, respectively (Figure S16). We then repeated our noise analysis for ATG measurements and observed that these perturbations led to down-weighting of the observations relating to FORS, SIMR and SKAG by $\sim 29\%$, 48% and 12% , respectively (Table S8). We similarly ran the multi-stage solutions for the new data set as per the unperturbed case. The results show that these perturbations caused changes of $+0.15$, $+0.28$ and $+0.07$ mm/yr in bias drift estimates of Jason-1, OSTM/Jason-2 and Jason-3, respectively, suggesting only slight sensitivity to the datum changes. However, our VLM estimates for the perturbed TGs changed significantly by $+1.53$, $+3.17$ and -0.12 mm/yr at FORS, SIMR and SKAG locations (as expected). The remainder of TG VLMs also experienced a shift with a weighted average of -0.08 mm/yr and WRMSE of 0.17 mm/yr (Figure S17). This experiment emphasizes the importance of detection and removal of large datum errors in providing reliable estimates of TG VLMs (see Text S2.4.1).

To examine possible effects of small spatial scale variability in our bias drifts on ASL estimates, we separately ran the multi-stage solutions for the southern and northern parts of the Baltic catchment. The resultant bias drifts from the sub-domains are within ± 0.06 mm/yr of those from the full domain (Table S9). The VLM differences between sub-domains and our full domain have a WRMSE of 0.12 mm/yr (Figure S18). We thus retrieved the same latitudinal pattern for ASL at TGs (see Figure S19). As expected, we saw slightly increased uncertainty (by a factor of ~ 1.5) for bias drift estimates from each sub-domain solution.

To inspect the impact of bias drift corrections on ASL estimates, we further compared our ASL at TGs with those where TG VLMs were derived with bias drifts constrained to be zero. The RMSE of ASL trends about the fitted polynomial is 0.88 mm/yr using TG VLMs from the zero-drift solution (only marginally higher than 0.87 mm/yr from the nominal solution). This experiment, however, suggests that discarding bias drift estimates would mostly affect the absolute magnitude of the ASL in the region - here we find that on average ASL is overestimated by ~ 0.3 mm/yr along the coasts (see Figures S20 and S21).

Finally, we repeated our multi-stage solutions using a priori VLMs taken from the ICE6G_D model with a nominal uncertainty of ± 1 mm/yr (Text S2.4.3). This led to a positive shift in our bias drift estimates ranging from $+0.71$ to $+0.80$ mm/yr, compared with the preferred solution (Figure S23). This is compatible with the difference between a priori TG VLMs derived from the GIA and GPS-Krig over the Fennoscandian region (a weighted average of -0.69 mm/yr and standard deviation of 0.71 mm/yr, noting the difference in sign is expected given the inverse relationship between bias drift and VLM).

4.5. Residuals and a Posteriori Analysis

To test if our Kalman filter operated effectively, we assessed the convergence between estimates of filtering and smoothing solutions. We postulated that convergence of bias drifts (e.g., after ~ 2.5 years as seen in Figure 7) is an indication that these estimates are separable from other temporally correlated parameters. We individually investigated the time series of our a posteriori estimates of cycle-by-cycle averaged “white plus AR1” noise on a per TG basis, location-specific TG VLM, ATG intercept pertaining to a TG/CP pair, and CP-specific ASL slope, ensuring appropriate decorrelation of parameters from the noise content in the multi-stage approach. We show the relevant outputs for a representative TG and CP in Figures S24–S27. We also checked the spatial coherence of ASL slopes over the region, and the deviations from a priori values (Figure S29). We then computed the weighted average “stack” of the “white plus AR1” residuals present in the ATG combinations of all TGs from our final solution to investigate the presence of any unmodeled time-variable signals. Figure 8 shows these residuals, which provide evidence of non-linear signals which we return to in the discussion.

We further evaluated the robustness of our results through a suite of a posteriori quality control analysis. First, we computed the cycle-by-cycle weighted average of differences between our TG VLM estimates and a priori GPS-Krig values. As expected, these were small and did not exceed our a priori “datum” uncertainty of ± 0.59 mm/yr (see Figure 3).

Second, we assessed the final estimates of ATG intercept that may have the potential to soak up VLM or bias drift signals. Note ATG intercepts were treated as time-variable parameters in the initial step of our multi-stage approach, and as time-fixed quantities in the subsequent solutions (see Section 2.3). We constructed the weighted average of ATG intercepts from Solution 3 on a per TG basis, after the removal of the median

Figure 8. Cycle-by-cycle weighted average stack of “white plus AR1” residuals of altimeter minus tide gauge (ATG) combinations over the region from the final solution of our multi-stage approach, with mission-specific root mean squared error (RMSE) annotated.

of individual intercept series. No significant residual trend remained, suggesting our ATG intercepts are effectively robust for subsequent analysis.

Lastly, we searched for evidence of spatially correlated common-mode errors in the estimated TG VLMs that would arise if there was imperfect modeling of bias drifts. We computed the cycle-by-cycle weighted average TG VLMs and monitored the stability over time. The results show a smooth transition between missions suggesting negligible decorrelation errors (Figure S28), and no significant step-change in bias drift estimates over any one mission lifespan. This adds confidence to our interpretation of the relatively high bias drift estimate for the TOPEX-side B mission. See Text S2.5 for more details.

5. Discussion

We have estimated, in the same solution, VLM trends and regional altimeter systematic errors using ALT, TG and GPS data. A serious challenge was the separation of mission-specific bias drifts and TG-specific VLMs in the presence of time-correlated noise. The inherent assumption of our solution is that the a priori VLMs interpolated from the GPS network are reasonable at most TG locations. This condition effectively sets a stable datum for the first solution and enables refinement of both bias drift and VLM in subsequent solutions.

In the context of the Baltic Sea as our case study, it is assumed that GIA is the main driver of VLM (e.g., Fjeldskaar, 1994; Lambeck et al., 1998; Milne et al., 2001). We expected, however, non-GIA factors also influence the ASL estimation from TG records in this region. GPS and GIA VLM trends at GPS locations reveal a median discrepancy of +0.39 mm/yr, and a standard deviation of 0.98 mm/yr, consistent with the findings of Johansson et al. (2002) and Lidberg et al. (2010). The important question is whether TG locations experience similar or greater local VLM variability that simply cannot be inferred by either interpolated GPS or GIA VLMs.

Our approach suggests that local variability is present, yielding VLM estimates at some TGs significantly different from interpolated GPS trends (Figure 3). For example, SASS, DAUG, SKUL and RODV TGs were estimated to have VLM differences of +1.26, +2.92, +4.08 and −4.53 mm/yr from the nearest GPS at the SASS, OJAR, LIMB and BUDP sites located 0.3, 14.1, 28.3 and 54.5 km away, respectively. These indicate the potential of localized variability, at least at some TGs, mostly in the central and southern regions (Figures 2b and S9b). An alternate hypothesis is that these anomalies arise from undocumented changes to individual RSL datums. However, the cycle-by-cycle weighted averages of ATG “white plus AR1” residuals associated with these gauges show no obvious shifts indicative of datum errors. Previous TG VLM estimates in the region by Kuo et al. (2004) and Santamaría-Gómez et al. (2014) differed from our preferred solution with weighted averages of the differences of +1.51 and +1.11 mm/yr and the WRMSE of 0.75 and 0.67 mm/yr, respectively. It is difficult to provide meaningful interpretation of these comparisons given significant

differences in time span considered, as well as different terrestrial reference frames and dissimilar altimetry products used. Our VLM estimates improved the spatial coherency of ASL and advocate that the use of interpolated GPS or GIA models is inadequate for accurately estimating ASL at TG locations.

We fitted a quadratic polynomial to the ASL derived using our VLMs versus those from GPS-Krig and GIA alternatives, and evaluated the sea-level variability by latitude. Our approach gives a $\sim 20\%$ reduction in RMSE of ASL variability (Figure 5). Our ASL estimates at TGs show good agreement with those from altimetry alone at CPs using our bias drifts and relative biases (Figure 6). We inferred a latitudinal dependence for ASL variability across the coasts that is consistent with the finding from the SLCCI altimetry data set (Madsen et al., 2019) and explained by climate-driven processes (Graham, 2004; Granskog et al., 2006; Hansson et al., 2011; Lehmann et al., 2011). We deduced an increasing rate of ASL by latitude, such that the weighted averages of our ASL trends in the southern and northern regions are $+3.67 \pm 0.44$ and $+5.23 \pm 0.46$ mm/yr, each slightly higher than the global average rate (e.g., $\sim +3.0 \pm 0.4$ mm/yr, Nerem et al., 2018).

Our evidence supports the presence of small but significant regional systematic errors in altimetry records with magnitudes that are larger than the mean estimates reported in global assessments (e.g., C. S. Watson et al., 2015), yet consistent with expectations based on orbit differences (Couhert et al., 2015; Fu & Haines, 2013). Our bias drifts appear robust as confirmed by the solution using a priori ICE6G_D VLMs, slightly shifting them in positive direction compatible with the discrepancy between GIA and GPS-interpolated values. Ignoring altimeter systematic errors hinders our ability to reach the regional accuracy goal of $\sim \pm 1$ mm/yr for VLM (Wöppelmann & Marcos, 2016), and $\sim \pm 0.5$ mm/yr for ASL (Ablain et al., 2015).

We investigated the implication of altimeter bias drifts for ASL estimates at CPs using altimetry alone, applying our inter- and intra-mission biases but not applying the bias drifts (Figure S22). Neglecting these small systematic errors led to an overestimation of the average magnitude of ASL by ~ 0.3 mm/yr. We reached the same conclusion with ASL at TGs, estimated by applying our VLMs to RSL trends, underscoring the importance of altimeter performance evaluation at regional scales (Figure S21).

It is well recognized that the detection and removal of often pervasive RSL datum errors are challenging when using TG data for any trend analysis (e.g., Nerem & Mitchum, 2002; Wöppelmann et al., 2008; C. S. Watson et al., 2015; Woodworth et al., 2017). Our sensitivity experiment suggests the effects on bias drifts are minimal when the datum errors are mitigated by automatic down-weighting of the ATG observations, except for large shifts (e.g., 50 mm) that were not manually detected. Our approach can detect TGs with potential datum issues as part of the solutions, which may flag further investigation and resolution of site-specific effects. This process enabled us to identify the GDANSK/NOWY PORT (GDAN) as a suspect TG (Figure S15, datum offset of ~ 55 mm at ~ 2013.7), confirmed by later differencing against the WLADYS-LAWOWO (WLAD) TG located 48 km away. Our tests show that we were able to notice considerable datum errors of +30 and +50 mm, but a small datum shift of +10 mm could not be easily detected (Section 4.4). This breakdown point raises the alternate hypothesis: TGs identified with anomalous VLM may alternatively (or in addition) suffer from unresolved changes to their RSL datums. The worst-case scenario is a small datum change placed at the midpoint time of the satellite era, e.g., a 15 mm datum shift at the midpoint time of 2006.45 would impart a linear trend error of 0.54 mm/yr over the full record. Therefore, datum instabilities may still contribute to what we recognized as VLM variability.

We assessed the sensitivity to a priori TG VLMs by our “mild” and “extreme” perturbation experiments. These suggest that over 80% of the magnitude of VLM perturbations are recovered if a priori estimates related to $\sim 15\%$ of the TGs were incorrect. The perturbations only partly affect bias drift estimates (changes of up to ~ -0.07 mm/yr and $\sim +0.32$ mm/yr for the mild and extreme scenarios respectively) and partially influence the VLM of the unperturbed gauges (Figure S14). Investigating the use of ATG trends outside our engine as a priori estimates of VLM highlights the importance of these nominal values in setting the datum and consequently our ability to resolve altimeter systematic errors. Keeping the a priori values independent of altimetry is therefore necessary, and thus we rely heavily on reasonable coverage of GPS-interpolated VLMs.

The assumption that “whole-of-system” bias drifts evolve linearly over the timespan of an individual mission is undoubtedly optimistic. Global analyses have shown non-linear effects with the TOPEX mission (e.g., C. S. Watson et al., 2015, and Beckley et al., 2017), and any comparison between competing orbit solutions shows a complex behavior in space and time (e.g., Couhert et al., 2015). Our solution investigates

the weighted averaged “stack” of residuals present in the ATG combinations and observes any unmodeled time-variable signals (Figure 8). The RMSE of the residuals (the summation of the estimated white and AR1 noise magnitudes) was in reasonable agreement with the RMSE of residuals in global studies (e.g., C. S. Watson et al., 2015). We detected some expression of non-linear signals particularly with the OSTM/Jason-2 mission. It is difficult to assess whether this time-variable signal is “real”; however, signals of similar magnitude are present in some in-situ validation time series used to determine the altimeter absolute bias (e.g., Haines et al., 2020; C. S. Watson et al., 2020). Analysis of orbit differences also suggests that the magnitude of this signal is plausible over spatial scales of a few thousand kilometers (e.g., Ablain et al., 2015).

A requirement of all Kalman-type implementations is an appropriate tuning of the filter, that is, defining the measurement noise and covariances as well as the state process noise and covariances used to construct the Kalman gains. This tuning is mainly dependent on the a priori assumptions built into the measurement equation and dynamic model. Our configurations are also reliant on the intrinsic characteristics of the study area. For example, the VLM field over Fennoscandia is relatively stable in time and therefore a tight process noise would be fine to estimate the quite linear VLM trends. Filter divergence occurs when the balance between TG-specific VLMs and mission-specific bias drifts is not achieved between and across space and time. We addressed this through the multi-stage decorrelation of parameters (Figure 1b), arriving at data-driven solutions within a flexible framework that is suitable to derive variables of interest through proper tuning of the filter. We thus enabled accurate determination of the state parameters (e.g., TG VLMs) in the presence of significant noise (e.g., typical RMSE of ATG observations of ~30 mm at a single CP).

The assumptions and configurations used here may not be appropriate in areas with different characteristics. For example, looser process noise is required for VLM trend estimates where significant non-linearity is present. We recognized anomalous geodetic sites using a threshold process noise that is dependent on the variability of the study VLM field. We offered a linear model to set up the relation between VLM and observations (Text S1.1), but this may not be suitable where extreme non-linear signals are present. As part of future work, we may dynamically change and adapt our process noise for intercepts related to potentially suspect TGs, or those that experience co- or post-seismic deformation, thus characterizing datum issues without undue influence on the mission-specific estimates.

The average reduction of ~34% in the ATG variance following the removal of seasonality highlights the presence of residual signal of oceanographic origin between the ALT and TG sampling locations. Any potential time-dependent non-seasonal contribution from this unresolved signal is quite difficult to quantify and is likely a significant limitation in any ATG-style approach for the determination of VLM, especially in regions with complex and dynamic ocean conditions.

6. Conclusions

We developed a Kalman-based methodology to adjust ATG, altimeter tandem/dual crossover, and GPS bedrock height observations to solve for localized vertical land motion and regionally coherent altimeter systematic errors (bias drifts) simultaneously. The key to our data-driven approach is the gradual decorrelation of unknowns over space and time within a multi-stage procedure, with appropriate treatment of time-invariant and time-variant noise content and spatiotemporal covariances in our observations.

We demonstrated the performance of the method in the Baltic Sea, where an abundance of TG and GPS sites exist, and ~24% of TGs have GPS sites in the vicinity of 10 km. Our method detected localized VLM at some TGs, particularly those situated in the southern region, that cannot be inferred by either interpolated GPS or GIA VLMs. This yields a more spatially coherent set of ASL estimates at TGs when computed using RSL plus our VLMs, with a ~20% reduction in RMSE of a quadratic polynomial fitted to the ASL per latitude, as compared to those using GPS or GIA based VLMs. Our ASL estimates at TGs are in good agreement with those inferred from altimetry alone at CPs using our bias drifts and relative biases applied to the ALT records. A comparison between our ASL at TGs and those computed using GIA VLMs also underscores the importance of non-GIA contributors for monitoring long-term sea-level rise.

Our estimates of “whole-of-system” altimeter systematic errors are significant (ranging from –4.6 mm/yr for TOPEX-side B to +2.6 mm/yr for OSTM/Jason-2), and considered plausible given mission specifications

and known spatially coherent differences between competing orbit solutions. The solution when the ICE-6G_D GIA was used as a priori VLM at TGs slightly shifted bias drifts in positive direction, consistent with the discrepancy between the GIA and GPS-interpolated VLMs. Comparing our solutions to those where bias drifts were constrained to zero reveals that the ASL could be overestimated by ~ 0.3 mm/yr across the region if the mission-specific errors were not considered.

Our robust approach could recover above 80% of VLMs at a few TGs ($\sim 15\%$) with perturbed records, and was also able to flag RSL datum errors of at least 30 mm (with a likely threshold of ~ 15 mm). Our method does however remain dependent on a “reasonable” set of a priori estimates of VLM as obtained from a GPS network to initially constrain the “datum” of estimated drifts. This emphasizes the ongoing need to install GPS directly at the TG or the nearest feasible location for optimal GPS coverage, as highlighted in the long-standing recommendation from the Intergovernmental Oceanographic Commission (Woodworth et al., 2016). Some evidence of non-linearity in bias drifts was observed (e.g., OSTM/Jason-2) however further work is required to assess the capability of capturing the likely time-variable evolution of altimeter errors.

For our test region, improved results would be anticipated using comprehensively retracked altimeter products that would more robustly deal with ice and land contamination (e.g., Gommenginger et al., 2011, Passaro et al., 2018, 2021), compared to the simplistic CP selection approach applied here. Our proposed method is likely applicable to other regional- and global-scale studies, but further investigation and potential modifications are required to estimate space- and time-varying bias drifts and crustal motions, particularly in high-dynamic areas. This technique could be further enhanced by including other missions (e.g., ERS-1/2, Envisat, and Sentinel-3A/B). Further investigation is required to evaluate the impact of residual differential oceanographic signals that likely exist between TG and the ALT CP sampling locations. Further work is also required to assess the merit of the technique for global studies with the inclusion of sites with non-linear behavior (e.g., due to earthquakes), that are typically excluded in similar studies (e.g., C. S. Watson et al., 2015).

Data Availability Statement

The altimeter and GPS data used in this study are publicly available through <https://github.com/remkos/rads> and <http://geodesy.unr.edu/>, respectively. Some TG data are available publicly and can be accessed from Meteorological and Hydrological Institute in Sweden at <https://www.smhi.se/data/oceanografi/ladda-ner-oceanografiska-observationer/#param=sealevelrh2000,stations=all>, Meteorological Institute in Finland at <https://en.ilmatieteenlaitos.fi/download-observations#!/>, Environment, Geology and Meteorology Center in Latvia at <https://www.meteo.lv/en/hidrologija-datu-meklesana/?&nid=919>. The TG data at RODVIG (RODV) location was obtained from https://resources.marine.copernicus.eu/?option=com_csw&view=details&product_id=INSITU_BAL_NRT_OBSERVATIONS_013_032. The remaining TG data are not publicly available and were provided on request by the data owners: Federal Maritime and Hydrographic Agency in Germany. Department of Marine Research of the Environmental Protection Agency in Lithuania, Institute of Meteorology and Water Management in Poland. Dynamic atmospheric Corrections are produced by CLS using the Mog2D model from Legos and distributed by Aviso+, with support from CNES (<https://www.aviso.altimetry.fr/>). The ICE6G_D GIA model is available through <http://www.atmosp.physics.utoronto.ca/~peltier/data.php>.

References

- Abblain, M., Cazenave, A., Larnicol, G., Balmaseda, M., Cipollini, P., Faugère, Y., et al. (2015). Improved sea level record over the satellite altimetry era (1993–2010) from the Climate Change Initiative project. *Ocean Science*, 11(1), 67–82. <https://doi.org/10.5194/os-11-67-2015>
- Altamimi, Z., Collilieux, X., & Métivier, L. (2011). ITRF2008: An improved solution of the international terrestrial reference frame. *Journal of Geodesy*, 85, 457–473. <https://doi.org/10.1007/s00190-011-0444-4>
- Beckley, B. D., Callahan, P. S., Hancock, D. W., Mitchum, G. T., & Ray, R. D. (2017). On the “cal-mode” correction to TOPEX satellite altimetry and its effect on the global mean sea level time series. *Journal of Geophysical Research: Oceans*, 122(11), 8371–8384. <https://doi.org/10.1002/2017jc013090>
- Benveniste, J., Birol, F., Calafat, F., Cazenave, A., Dieng, H., Gouzenes, Y., et al. (2020). Coastal sea level anomalies and associated trends from Jason satellite altimetry over 2002–2018. *Scientific Data*, 7(1), 357. <https://doi.org/10.1038/s41597-020-00694-w>
- Blewitt, G., Hammond, W. C., & Kreemer, C. (2018). Harnessing the GPS data explosion for interdisciplinary science. *Eos*, 99. <https://doi.org/10.1029/2018EO104623>

Acknowledgments

This research was supported under the Australian Research Council’s Special Research Initiative for Antarctic Gateway Partnership (Project ID SR140300001). This work has benefitted from the Bass Strait altimeter validation facility which is supported by Australia’s Integrated Marine Observing System (IMOS) - IMOS is enabled by the National Collaborative Research Infrastructure Strategy (NCRIS). IMOS is operated by a consortium of institutions as an unincorporated joint venture, with the University of Tasmania as Lead Agent. Aspects of this work were also supported by the Australian Research Council Discovery Project DP150100615. We acknowledge the TPAC High Performance Computing facilities used for computations. Discussions with Dr. Benoit Legresy and Prof. Richard Coleman while undertaking this work are appreciated. The authors thank Prof. Don P. Chambers (Editor), Dr. Alvaro Santamaría-Gómez (reviewer), and an anonymous reviewer for their constructive comments that helped to improve this manuscript.

- Bos, M. S., Fernandes, R. M. S., Williams, S. D. P., & Bastos, L. (2013). Fast error analysis of continuous GNSS Observations with missing data. *Journal of Geodesy*, 87(4), 351–360. <https://doi.org/10.1007/s00190-012-0605-0>
- Bouin, M. N., & Wöppelmann, G. (2010). Land motion estimates from GPS at tide gauges: A geophysical evaluation. *Geophysical Journal International*, 180(1), 193–209. <https://doi.org/10.1111/j.1365-246x.2009.04411.x>
- Buttkus, B. (2000). Spectral analysis and filter theory in applied geophysics (1st ed.). Springer-Verlag Berlin Heidelberg. <http://doi.org/10.1007/978-3-642-57016-2>
- Carrere, L., Lyard, F., Cancet, M., Guillot, A., & Picot, N. (2016). FES 2014, a new tidal model: Validation results and perspectives for improvements. In Proceedings at ESA Living Planet Conference, Prague, Czech Republic, European Space Agency.
- Cazenave, A., Dieng, H.-B., Meyssignac, B., Schuckmann, K. V., Decharme, B., & Berthier, E. (2014). The rate of sea-level rise. *Nature Climate Change*, 4(5), 358–361. <https://doi.org/10.1038/nclimate2159>
- Cazenave, A., Dominh, K., Ponchaut, F., Soudarin, L., Cretaux, J. F., & Provost, C. L. (1999). Sea level changes from Topex-Poseidon altimetry and tide gauges, and vertical crustal motions from DORIS. *Geophysical Research Letters*, 26(14), 2077–2080. <https://doi.org/10.1029/1999gl900472>
- Chiles, J.-P., & Delfiner, P. (2012). *Geostatistics: Modeling spatial uncertainty* (2nd ed.). John Wiley & Sons, Inc. <http://doi.org/10.1002/9781118136188>
- Church, J. A., Clark, P. U., Cazenave, A., Gregory, J. M., Jevrejeva, S., Levermann, A., et al. (2013). Sea-level rise by 2100. *Science*, 342(6165), 1445. <https://doi.org/10.1126/science.1242154>
- Church, J. A., & White, N. J. (2011). Sea-level rise from the late 19th to the early 21st century. *Surveys in Geophysics*, 32, 585–602. <https://doi.org/10.1007/s10712-011-9119-1>
- Codiga, D. L. (2011). Unified Tidal Analysis and Prediction Using the UTide Matlab Functions. Technical Report 2011-01, Graduate School of Oceanography, University of Rhode Island, Narragansett, RI. 59 pp. Retrieved from <http://www.po.gso.uri.edu/~codiga/utide/utide.htm>
- Couhert, A., Cerri, L., Legeais, J. F., Ablain, M., Zelensky, N. P., Haines, B. J., et al. (2015). Towards the 1 mm/y stability of the radial orbit error at regional scales. *Advances in Space Research*, 55(1), 2–23. <https://doi.org/10.1016/j.asr.2014.06.041>
- Desai, S., Wahr, J., & Beckley, B. (2015). Revisiting the pole tide for and from satellite altimetry. *Journal of Geodesy*, 89(12), 1233–1243. <https://doi.org/10.1007/s00190-015-0848-7>
- Featherstone, W. E., Penna, N. T., Filmer, M. S., & Williams, S. D. P. (2015). Nonlinear subsidence at Fremantle, a long-recording tide gauge in the Southern Hemisphere. *Journal of Geophysical Research: Oceans*, 120(10), 7004–7014. <https://doi.org/10.1002/2015jc011295>
- Fjeldskaar, W. (1994). Viscosity and thickness of the asthenosphere detected from the Fennoscandian uplift. *Earth and Planetary Science Letters*, 126(4), 399–410. [https://doi.org/10.1016/0012-821x\(94\)90120-1](https://doi.org/10.1016/0012-821x(94)90120-1)
- Fu, L.-L., & Haines, B. J. (2013). The challenges in long-term altimetry calibration for addressing the problem of global sea level change. *Advances in Space Research*, 51(8), 1284–1300. <https://doi.org/10.1016/j.asr.2012.06.005>
- Gommenginger, C., Thibaut, P., Fenoglio-Marc, L., Quartly, G., Deng, X., Gómez-Enrí, J., et al. (2011). Retracking altimeter waveforms near the coasts. In S. Vignudelli, et al. (Eds.), *Coastal altimetry* (pp. 6–101). Berlin: Springer.
- Graham, L. P. (2004). Climate change effects on river flow to the Baltic Sea. *Ambio*, 33(4–5), 235–241. <https://doi.org/10.1579/0044-7447-33.4.235>
- Granskog, M., Kaartokallio, H., Kuosa, H., Thomas, D. N., & Vainio, J. (2006). Sea ice in the Baltic Sea: A review. *Estuarine, Coastal and Shelf Science*, 70, 145–160. <https://doi.org/10.1016/j.ecss.2006.06.001>
- Grewal, M. S., & Andrews, A. P. (2008). *Kalman filtering: Theory and practice using MATLAB®* (3rd ed.). WILEY. <http://doi.org/10.1002/9780470377819>
- Haines, B., Desai, S. D., Kubitschek, D., & Leben, R. R. (2020). A brief history of the Harvest experiment: 1989–2019. *Advances in Space Research*. <https://doi.org/10.1016/j.asr.2020.08.013>
- Hammond, W. C., Blewitt, G., & Kreemer, C. (2016). GPS imaging of vertical land motion in California and Nevada: Implications for Sierra Nevada uplift. *Journal of Geophysical Research: Solid Earth*, 121(10), 7681–7703. <https://doi.org/10.1002/2016jb013458>
- Han, W., Stammer, D., Thompson, P., Ezer, T., Palanisamy, H., Zhang, X., et al. (2019). Impacts of basin-scale climate modes on coastal sea level: A review. *Surveys in Geophysics*, 40(6), 1493–1541. <https://doi.org/10.1007/s10712-019-09562-8>
- Hansson, D., Eriksson, C., Omstedt, A., & Chen, D. (2011). Reconstruction of river runoff to the Baltic Sea, AD 1500–1995. *International Journal of Climatology*, 31(5), 696–703. <https://doi.org/10.1002/joc.2097>
- Hay, C. C., Morrow, E., Kopp, R. E., & Mitrovica, J. X. (2015). Probabilistic reanalysis of twentieth-century sea-level rise. *Nature*, 517, 481–484. <https://doi.org/10.1038/nature14093>
- Hill, E. M., Davis, J. L., Tamisiea, M. E., & Lidberg, M. (2010). Combination of geodetic observations and models for glacial isostatic adjustment fields in Fennoscandia. *Journal of Geophysical Research*, 115(B7), B07403. <https://doi.org/10.1029/2009jb006967>
- Hünicke, B., & Zorita, E. (2008). Trends in the amplitude of Baltic Sea level annual cycle. *Tellus Series A: Dynamic Meteorology and Oceanography*, 60 A(1), 154–164. <https://doi.org/10.1111/j.1600-0870.2007.00277.x>
- Johansson, J. M., Davis, J. L., Scherneck, H.-G., Milne, G. A., Vermeer, M., Mitrovica, J. X., et al. (2002). Continuous GPS measurements of postglacial adjustment in Fennoscandia I. Geodetic results. *Journal of Geophysical Research*, 107(B8), ETG3-1–ETG3-27. <https://doi.org/10.1029/2001jb000400>
- King, M. A., Keshin, M., Whitehouse, P. L., Thomas, I. D., Milne, G., & Riva, R. E. M. (2012). Regional biases in absolute sea-level estimates from tide gauge data due to residual unmodeled vertical land movement. *Geophysical Research Letters*, 39(14), L14604. <https://doi.org/10.1029/2012gl052348>
- Kleinherenbrink, M., Riva, R., & Frederikse, T. (2018). A comparison of methods to estimate vertical land motion trends from GNSS and altimetry at tide gauge stations. *Ocean Science*, 14(2), 187–204. <https://doi.org/10.5194/os-14-187-2018>
- Kuo, C. Y., Shum, C. K., Braun, A., & Mitrovica, J. X. (2004). Vertical crustal motion determined by satellite altimetry and tide gauge data in Fennoscandia. *Geophysical Research Letters*, 31(1), 1–4. <https://doi.org/10.1029/2003gl019106>
- Lambeck, K., Smither, C., & Ekman, M. (1998). Tests of glacial rebound models for Fennoscandia based on instrumented sea- and lake-level records. *Geophysical Journal International*, 135(2), 375–387. <https://doi.org/10.1046/j.1365-246x.1998.00643.x>
- Lehmann, A., Getzlaff, K., & Harlaß, J. (2011). Detailed assessment of climate variability in the Baltic Sea area for the period 1958 to 2009. *Climate Research*, 46(2), 185–196. <https://doi.org/10.3354/cr00876>
- Lidberg, M., Johansson, J. M., Scherneck, H.-G., & Milne, G. A. (2010). Recent results based on continuous GPS observations of the GIA process in Fennoscandia from BIFROST. *Journal of Geodynamics*, 50(1), 8–18. <https://doi.org/10.1016/j.jog.2009.11.010>
- Madsen, K. S., Hoyer, J. L., Suursaar, Ü., She, J., & Knudsen, P. (2019). Sea level trends and variability of the baltic sea from 2D statistical reconstruction and altimetry. *Frontiers of Earth Science*, 7, 243. <https://doi.org/10.3389/feart.2019.00243>

- Milne, G. A., Davis, J. L., Mitrovica Jerry, X., Scherneck, H.-G., Johansson, J. M., Vermeer, M., & Koivula, H. (2001). Space-geodetic constraints on glacial isostatic adjustment in Fennoscandia. *Science*, 291(5512), 2381–2385. <https://doi.org/10.1126/science.1057022>
- Montero, J. -M., Fernández-Avilés, G., & Mateu, J. (2015). *Spatial and spatio-temporal geostatistical modeling and Kriging*. John Wiley & Sons, Ltd. <http://doi.org/10.1002/9781118762387>
- Nerem, R. S., Beckley, B. D., Fasullo, J. T., Hamlington, B. D., Masters, D., & Mitchum, G. T. (2018). Climate-change-driven accelerated sea-level rise detected in the altimeter era. *Proceedings of the National Academy of Sciences of the United States of America*, 115(9), 2022–2025. <https://doi.org/10.1073/pnas.1717312115>
- Nerem, R. S., & Mitchum, G. T. (2002). Estimates of vertical crustal motion derived from differences of TOPEX/POSEIDON and tide gauge sea level measurements. *Geophysical Research Letters*, 29(19), 41–44. <https://doi.org/10.1029/2002gl015037>
- Neumann, B., Vafeidis, A. T., Zimmermann, J., & Nicholls, R. J. (2015). Future coastal population growth and exposure to sea-level rise and coastal flooding - A global assessment. *PLoS ONE*, 10(3). <https://doi.org/10.1371/journal.pone.0118571>
- Oelsmann, J., Passaro, M., Dettmering, D., Schwatke, C., Sanchez, L., & Seitz, F. (2021). The zone of influence: Matching sea level variability from coastal altimetry and tide gauges for vertical land motion estimation. *Ocean Science*, 17(1), 35–57. <https://doi.org/10.5194/os-17-35-2021>
- Ostanciaux, É., Husson, L., Choblet, G., Robin, C., & Pedoja, K. (2012). Present-day trends of vertical ground motion along the coast lines. *Earth-Science Reviews*, 110(1–4), 74–92. <https://doi.org/10.1016/j.earscirev.2011.10.004>
- Passaro, M., Müller, F., Dettmering, D., Abulaitjiang, A., Rautiainen, L., Scarrott, R. G., et al. (2021). Baltic SEAL: Product handbook, Version 1.1. Report delivered under the Baltic SEAL project (ESA contract no. 000126590/19/I/BG). <http://doi.org/10.5270/esa.BalticSEAL.PH1.1>
- Passaro, M., Rose, S. K., Andersen, O. B., Boergens, E., Calafat, F. M., Dettmering, D., & Benveniste, J. (2018). ALES+: Adapting a homogeneous ocean retracker for satellite altimetry to sea ice leads, coastal and inland waters. *Remote Sensing of Environment*, 211, 456–471. <https://doi.org/10.1016/j.rse.2018.02.074>
- Peltier, W. R., Argus, D. F., & Drummond, R. (2018). Comment on “An assessment of the ICE-6G_C (VM5a) Glacial isostatic adjustment model” by Purcell et al. *Journal of Geophysical Research: Solid Earth*, 123(2), 2019–2028. <https://doi.org/10.1002/2016jb013844>
- Pfeffer, J., & Allemand, P. (2016). The key role of vertical land motions in coastal sea level variations: A global synthesis of multisatellite altimetry, tide gauge data and GPS measurements. *Earth and Planetary Science Letters*, 439, 39–47. <https://doi.org/10.1016/j.epsl.2016.01.027>
- Pfeffer, J., Spada, G., Mémin, A., Boy, J. P., & Allemand, P. (2017). Decoding the origins of vertical land motions observed today at coasts. *Geophysical Journal International*, 210(1), 148–165. <https://doi.org/10.1093/gji/ggx142>
- Santamaría-Gómez, A., Bouin, M.-N., Collilieux, X., & Wöppelmann, G. (2011). Correlated errors in GPS position time series: Implications for velocity estimates. *Journal of Geophysical Research*, 116(1), B01405. <https://doi.org/10.1029/2010jb007701>
- Santamaría-Gómez, A., Gravelle, M., & Wöppelmann, G. (2014). Long-term vertical land motion from double-differenced tide gauge and satellite altimetry data. *Journal of Geodesy*, 88(3), 207–222. <https://doi.org/10.1007/s00190-013-0677-5>
- Scharroo, R., Leuliette, E. W., Lillibridge, J. L., Byrne, D., Naeije, M. C., & Mitchum, G. T. (2013). RADS: Consistent multi-mission products, in *Proceedings Symposium on 20 Years of Progress in Radar Altimetry*, European Space Agency, Special Publication SP-710.
- Spada, G., & Galassi, G. (2012). New estimates of secular sea level rise from tide gauge data and GIA modelling. *Geophysical Journal International*, 191(3), 1067–1094. <https://doi.org/10.1111/j.1365-246X.2012.05663.x>
- Watson, C. S., Legresy, B., Beardsley, J., Zhou, A., & King, M. A. (2020). Altimeter validation results from the Bass Strait validation facility, Australia. Ocean Surface Topography Science Team Meeting 2020, Virtual Forum. Retrieved from https://ostst.avisio.altimetry.fr/fileadmin/user_upload/tx_ausysclseminar/files/OSTST2020_Absolute_Bias_Watson_et_al.pdf
- Watson, C. S., Legresy, B., & King, M. A. (2019). On the uncertainty associated with validating the global mean sea level climate record. *Advances in Space Research*. <https://doi.org/10.1016/j.asr.2019.09.053>
- Watson, C. S., White, N. J., Church, J. A., King, M. A., Burgette, R. J., & Legresy, B. (2015). Unabated global mean sea-level rise over the satellite altimeter era. *Nature Climate Change*, 5(6), 565–568. <https://doi.org/10.1038/nclimate2635>
- Watson, P. J. (2019). An Assessment of the utility of satellite altimetry and tide gauge Data (ALT-TG) as a proxy for estimating vertical land motion. *Journal of Coastal Research*, 35(6), 1131–1144. <https://doi.org/10.2112/jcoastres-d-19-00031.1>
- Williams, S. D. P., Bock, Y., Fang, P., Jamason, P., Nikolaidis, R. M., Prawirodirdjo, L., et al. (2004). Error analysis of continuous GPS position time series. *Journal of Geophysical Research*, 109(3), 1–19. <https://doi.org/10.1029/2003jb002741>
- Woodworth, P. L., et al. (Eds.), (2016). *Manual on Sea-level Measurements and Interpretation*, Volume V: Radar Gauges. Paris, Intergovernmental Oceanographic Commission of UNESCO. 104 pp & Supplement 126 pp. (IOC Manuals and Guides No.14, vol. V; JCOMM Technical Report No. 89; (English).
- Woodworth, P. L., Gehrels, W. R., & Nerem, R. S. (2011). Nineteenth and twentieth century changes in sea level. *Oceanography*, 24(2), 80–93. <https://doi.org/10.5670/oceanog.2011.29>
- Woodworth, P. L., Wöppelmann, G., Marcos, M., Gravelle, M., & Bingley, R. M. (2017). Why we must tie satellite positioning to tide gauge data. *Eos*, 98. <https://doi.org/10.1029/2017EO064037>
- Woolf, D. K., Shaw, A. G. P., & Tsimplis, M. N. (2003). The influence of the North Atlantic Oscillation on sea-level variability in the North Atlantic region. *Journal of Atmospheric and Oceanic Technology*, 20(4), 145–167. <https://doi.org/10.1029/2002JO001633>
- Wöppelmann, G., Letetrel, C., Santamaría, A., Bouin, M.-N., Collilieux, X., Altamimi, Z., et al. (2009). Rates of sea-level change over the past century in a geocentric reference frame. *Geophysical Research Letters*, 36, L12607. <https://doi.org/10.1029/2009gl038720>
- Wöppelmann, G., & Marcos, M. (2016). Vertical land motion as a key to understanding sea level change and variability. *Reviews of Geophysics*, 54(1), 64–92. <https://doi.org/10.1002/2015rg000502>
- Wöppelmann, G., Pouvreau, N., Coulomb, A., Simon, B., & Woodworth, P. L. (2008). Tide gauge datum continuity at Brest since 1711: France's longest sea-level record. *Geophysical Research Letters*, 35, L22605. <https://doi.org/10.1029/2008GL035783>

Reference From the Supporting Information

- Nerem, R. S., Chambers, D. P., Choe, C., & Mitchum, G. T. (2010). Estimating mean sea level change from the TOPEX and Jason altimeter missions. *Marine Geodesy*, 33(1), 435–446. <https://doi.org/10.1080/01490419.2010.491031>