



Short note: extracellular export and consumption of glucose in Antarctic sea ice

Fraser Kennedy¹ · Andrew McMinn^{1,2} · Andrew Martin¹

Received: 3 November 2021 / Revised: 21 February 2022 / Accepted: 23 February 2022 / Published online: 11 March 2022
© The Author(s) 2022

Abstract

Extracellular carbohydrate production is widespread in sea ice microbial communities, being produced by both algae and bacteria. Under stressful conditions, including nutrient limitation and high light, cells may export excess fixed carbon as glucose. Glucose microsensors were used to measure extracellular glucose exudation and consumption in a sea ice algal community. Glucose export increased with increasing irradiance between 15 and 512 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. This export correlated with declining F_vF_m values and increasing NPQ values, implying that glucose export resulted from exposure to above optimal irradiances. Glucose concentrations in samples treated with DCMU to block photosynthesis, declined at all irradiances. Bacterial consumption of glucose was between 6 and 34% of extracellular export per hour. There have been very few measurements of DOC/glucose in sea ice and the data presented here make an important contribution to our understanding of sea ice microbial processes.

Keywords Glucose · Antarctic microsensor · Sea ice algae

Introduction

Polar sea ice contains high biomass, specifically diverse microbial communities that make a major contribution to the annual primary productivity of perennially ice-covered polar seas (Arrigo 2014). Biomass is typically dominated by diatom communities although other protists, bacteria, Archaea and viruses are also abundant (McMinn et al. 2020). Even though temperatures and under-ice irradiances are low, productivity rates are often high and comparable to productive communities elsewhere (Lizotte 2001; McMinn et al. 2012; van Leeuwe et al. 2018).

Extracellular carbohydrate production is widespread in sea ice microbial communities, being produced by both algae and bacteria (Ugalde et al. 2014). One class of carbohydrates, extracellular polymeric substances (EPS), has been shown to have multiple roles that include cell motility, substrate attachment and protection from grazing (Underwood

and Kromkamp 1999; Staats et al. 2000; Cook et al. 2007). Much of this EPS is composed of polysaccharides, uronic acids, and sulphated sugars (Underwood and Paterson 2003; Bellinger et al. 2009; Oakes et al. 2010; Rivaro et al. 2021). Microalgal dissolved organic carbon (DOC) production and export are fundamental characteristics that underpin the functioning of microbial food webs (Martin et al. 2011, 2012). After export from the microalgae, the DOC is rapidly taken up by both heterotrophic bacteria and the microalgae. It has been shown that most microalgae have the ability to directly take up glucose (Leles et al. 2019). Extracellular release of DOC from microalgae is mostly by either cell lysis, resulting from grazing or viral attack, or photosynthetic overflow, i.e. where photosynthetic carbon fixation exceeds that required for balanced macromolecular synthesis and the resulting excess photosynthate is released directly into the water column (Smith and Underwood 2000). The availability of an organic substrate is essential for heterotrophic bacterial growth and the effective functioning of the microbial loop (Azam et al. 1983). There is consequently often a close and coupled association between autotrophic producers, such as diatoms, and bacterial consumers (Fenchel 2008; Martin et al. 2012). The composition of DOC exudates from microalgae is dominated by glucose, which often comprises more than 70% of the total (Hama and Yanagi 2001; Haas

✉ Andrew McMinn
Andrew.mcminn@utas.edu.au

¹ Institute for Marine and Antarctic Studies, University of Tasmania, Private Bag 129, Hobart, TAS 7001, Australia

² College of Marine Life Sciences, Ocean University of China, Qingdao 266003, China

and Wild 2010; Underwood et al. 2010). The causes and drivers of extracellular DOC export (through photosynthetic overflow) are not well understood, but any process that leads to a reduction in growth, such as nutrient limitation or excessive light (Staats et al. 1999; Underwood and Paterson 2003; Cook et al. 2004, 2007), is likely to contribute. Extracellular DOC exudation in culture, for instance, is mostly associated with the low nutrient, stationary phase of growth rather than the exponential phase (Myklesstad 1977). Under ice nutrients are usually abundant and non-limiting during spring in McMurdo Sound (McMinn et al. 1999) and so growth limitation is most likely to result from insufficient light.

Whilst a range of organic carbon compound categories (e.g. lipids, proteins and carbohydrates) have previously been quantified in the EPS of sea ice (Palmisano and Sullivan 1985; Aslam et al. 2012; Ugalde et al. 2014), glucose itself was first identified as an important component of sea ice habitats by Underwood et al. (2010). Extracellular glucose production has since been documented by Aslam et al. (2016). Here we examine extracellular glucose exudation and consumption in sea ice exposed to a light gradient. We hypothesise that excessive irradiance, i.e. light levels high enough to cause photoinhibition, will lead to an increase in glucose exudation.

Materials and methods

Three replicate sea ice cores, within one metre of each other, were collected from Cape Evans, Antarctica (77° 37' 99" S, 166° 23' 99" E), by SIPRE corer on 12 November 2019.

The bottom 5 mm of each core was shaved into 500 ml of pre-chilled filtered sea water and stored in the dark at ~0 °C for 20 min for the cells to acclimate. This approach resulted in a drop in salinity of less than 2%. Chlorophyll *a* analysis of the sea ice community followed McMinn et al. (2007). Relative species analysis was based on a count of 400 cells using a Zeiss Axioskop microscope at 400×.

Subsamples (5 ml) from each ice core were placed in 3 ml cuvettes and exposed to light levels of 15, 68, 227, and 512 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ for four minutes for the glucose concentrations to be measured. This time period was considered sufficient to measure the change in glucose concentration over time (i.e. extracellular glucose production/consumption) but insufficient for algal and bacterial biomass to change. Glucose measurements were made every 1 s. The control sample, which contained the same samples but with the sea ice algae removed, was also measured at each irradiance. DCMU (concentration 20 mM) was added to an additional set of samples from the same cores and measured at the same range of irradiances. All incubations and measurements occurred at 0 °C.

Glucose measurements

A glucose biosensor (Pinnacle Technology Lawrence KS, USA) was used to measure net glucose exudation following McMinn and Lee (2018). This biosensor uses the oxidase enzyme to reduce the glucose molecules and produce hydrogen peroxide as a by-product. The hydrogen peroxide is reduced on a platinum electrode to produce electrons that are then detected on an Ag/AgCl reference electrode (McMinn and Lee 2018). As relatively high glucose concentrations are required for detection ($> 2\text{--}5 \mu\text{M L}^{-1}$), it was important to use relatively high biomass sea ice samples. The output of each biosensor (picoamps) was calibrated against a standard curve (D-glucose standard, Sigma Aldrich, St. Louis, MO, USA), composed of glucose solutions at concentrations of 0.125, 0.25, 0.375, 1.375 and 2.375 mM L^{-1} . This standard curve was used to convert the raw millivolt signal of the biosensors to actual glucose concentrations. The biosensors have a 90% response time of ~4 s (McMinn and Lee 2018).

The glucose biosensor was mounted onto a motorized micromanipulator (Unisense A/S, Aarhus, Denmark) and connected to a field multimeter (Unisense, A/S, Aarhus, Denmark); the raw data were converted into mg glucose L^{-1} using the relationship from the standard curve.

To obtain total glucose exudation, the glucose consumption in the dark (the control) was added to the glucose production in the light. Glucose consumption in the dark represents heterotrophic uptake of glucose by both algae and bacteria. The exudation rate was measured by determining the change in glucose concentration over two minutes.

Exudation rates were normalized to chlorophyll *a* (chl *a*) concentrations and time [$\mu\text{M glucose (mg chl } a)^{-1} \text{h}^{-1}$]. Chlorophyll *a* concentration of each replicate was measured by fluorometer (Turner 10AU, KA), using the acidification method, noting that the sea ice algal community was the same in each replicate.

Pulse amplification-modulated fluorometry (PAM)

Chlorophyll *a* fluorescence of photosystem II was measured using a pulse amplitude-modulated fluorometer (Water-PAM, Walz, Effeltrich, Germany) with an internal actinic light source centred on 660 nm. All samples were dark acclimated for 30 min prior to measurement.

A more complete description of the PAM methods used is described in Kennedy et al. (2020). Fluorescence induction curves, which were used to examine the effect of increasing irradiance on PSII reaction centre kinetics and energy dissipation mechanisms, were obtained under

software control (WinControl 3; Walz). Cells were first dark adapted for 30 min and then exposed to a brief pulse of far-red illumination (wavelength > 680 nm) for 5 s to oxidize PSI and the electron transport chain to gain an estimate of F_0 . Induction curves were initiated by first determining F_v/F_m by a saturation pulse, followed by a delay of 30 s before actinic illumination (either 15, 68, 227 or 516 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) was turned on. Additional saturation pulses were performed at the onset of actinic illumination and every 20 s thereafter until the cessation of actinic light four minutes later. The final saturation pulse at four minutes, with actinic light turned off, was used to determine the final F_v/F_m value and non-photochemical quenching (NPQ). NPQ represents the fraction of heat that is dissipated via regulated photoprotective mechanisms such as the xanthophyll cycle and is derived from:

$$\text{NPQ} = (F_m - F'_m) / F'_m \text{ (Genty et al. 1989)}$$

Statistics

Pearson correlations were performed to observe relationships between glucose exudation, F_v/F_m and NPQ using the software package Rstudio [RStudio team (2021), version 1.4.1106]. Data were checked for normality using the Shapiro–Wilk test.

Results

Sea ice at Cape Evans on 12 November 2019 was 1.25 m thick without a snow cover. Chlorophyll *a* concentration was $74.9 \pm 6.9 \text{ mg chl } a \text{ m}^{-2}$ and the ice algal community was dominated by *Nitzschia stellata* (52%), *Berkelaya adeliense* (36%) and *Navicula glaciei* (7%). Diatoms comprised > 99% of the protists.

Glucose measurements

Independent glucose measurements were made at each of the four light treatments (15, 68, 227, 512 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$). At each irradiance, additional samples were used to measure a control (no added sea ice algae) and samples treated with DCMU, to inhibit photosynthetic activity. At all light levels, glucose concentrations in the control samples were beneath the level of detection. Glucose concentrations across all other samples ranged from 0.012 to 0.167 mg L^{-1} . Net extracellular glucose export rates were 0.638 ± 1.062 , 0.844 ± 0.022 , 13.472 ± 11.374 and $9.942 \pm 1.282 \mu\text{M glucose (mg chl } a)^{-1} \text{ h}^{-1}$ at 15, 68, 227 and 512 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, respectively (Table 1). Samples treated with DCMU experienced a drop in glucose concentrations, presumably mostly due to bacterial consumption. Glucose consumption rates were 0.283 ± 0.112 , 0.799 ± 0.325 and $0.766 \pm 0.0005 \mu\text{M glucose (mg chl } a)^{-1} \text{ h}^{-1}$ at 68, 227 and 512 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, respectively (Table 1). Electrode response in the 15 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ sample was unstable and so no consumption rate could be measured. Drop in glucose concentrations in the dark (bacterial consumption) was equivalent to ~34, 6 and 8% of extracellular glucose production per hour in the light; these are equivalent to glucose pool turnover times of 2.9, 17 and 13 h^{-1} at 68, 227 and 512 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, respectively, although the large standard deviations for some of the measurements means these values are indicative only. There was a moderate negative correlation between extracellular glucose export and bacterial uptake ($r^2 = 0.68$, $p = 0.001$).

Pulse-amplitude modulation (PAM) measurements

F_v/F_m values at the beginning of the inductions curves at the four irradiances were between 0.331 ± 0.156 and 0.476 ± 0.019 (Table 1). However, at the end, after four minutes of actinic light exposure these had dropped from 0.303 ± 0.082 at 15 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ to 0.206 ± 0.120 ,

Table 1 Extracellular glucose export and consumption

Irradiance	Glucose export rate	Glucose consumption rate	Turnover time	F_v/F_m (start)	F_v/F_m (end)	% decline	NPQ
15	0.115 ± 0.193			0.476 ± 0.019	0.303 ± 0.082	36	0.253 ± 0.027
68	0.152 ± 0.004	0.051 ± 0.021	2.9	0.331 ± 0.156	0.206 ± 0.120	38	0.355 ± 0.248
227	2.427 ± 2.068	0.144 ± 0.059	17	0.371 ± 0.077	0.025 ± 0.031	93	0.432 ± 0.020
512	1.791 ± 0.233	0.138 ± 0.0001	13	0.471 ± 0.035	0.019 ± 0.018	96	0.779 ± 0.293

Irradiance ($\mu\text{mol photons m}^{-2} \text{s}^{-1}$); glucose export rate [i.e. net extracellular glucose export rate, $\text{mg glucose (mg chl } a)^{-1} \text{ h}^{-1}$]; glucose consumption rate (i.e. net reduction in the amount of exuded algal glucose by bacterial/algal glucose consumption) [$\text{mg glucose (mg chl } a)^{-1} \text{ h}^{-1}$]; turnover time (time to consume glucose pool h^{-1}); F_v/F_m (start) is the F_v/F_m value (dimensionless ratio) at the start of the induction curve after 30-min dark acclimation. F_v/F_m (end) is the F_v/F_m value (dimensionless ratio) at the end of the 4-min induction curve; % decline is the percentage drop in F_v/F_m values between the beginning and end of the induction curve

NPQ non-photosynthetic quenching (dimensionless)

Errors are standard deviations based on three replicates

0.025 ± 0.031 and 0.019 ± 0.018 at 68, 227 and 512 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, respectively. Over the same time interval, NPQ values increased from 0.253 ± 0.027 at 15 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ to 0.430 ± 0.298 , 0.696 ± 0.432 and 0.779 ± 0.293 at 68, 227 and 512 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, respectively (Table 1). There was a moderate correlation between extracellular glucose export and % decline in F_v/F_m ($r^2 = 0.75$, $p = 0.00026$) and a weak correlation between extracellular glucose export and maximum NPQ ($r^2 = 0.39$, $p = 0.031$).

Discussion

In this study, net extracellular glucose export in a sea ice community from McMurdo Sound increased with increasing irradiance, with a major increase between 68 and 227 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. DOC production and exudation in marine biofilms, including sea ice, have been well documented and has usually been found to be associated with high light levels and/or nutrient limitation (Staats et al. 1999; Underwood and Paterson 2003; Cook et al. 2004, 2007; Underwood et al. 2013; Ugalde et al. 2014). Although nutrient levels were not measured, other studies have found Antarctic under-ice nutrient concentrations during spring to be high and non-limiting (McMinn et al. 1995, 1999; Cummings et al. 2019). Most studies have shown that sea ice communities reach light saturation, i.e. have E_k values, at irradiances of less than 200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, with inhibition starting at $< 300 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ (McMinn et al. 2012; Sorrell et al. 2021). Similarly, in situ communities exposed to ambient irradiances have F'_v/F_m values of < 0.1 , compared with night-time values of > 0.4 (McMinn et al. 2003). Most studies actually show E_k values less than 100 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ (McMinn et al. 2003, 2007, 2010; Sorrell et al. 2021). In this study also, there was a sharp drop in final F_v/F_m and an increase in NPQ values, between irradiances of 68 and 227 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. NPQ is a protective mechanism that is upregulated to divert energy from excess irradiance to prevent cellular damage. In this study NPQ increased at all irradiances but was much greater after exposure to higher irradiances. There is a weak correlation between these increased NPQ values and glucose export ($r^2 = 0.370$). As under-ice inorganic nutrient levels in spring in McMurdo Sound are consistently high (Arrigo 2014; Cummings et al. 2019), this relationship implies that it is likely that the above optimal irradiances alone caused the increase in DOC exudation (glucose) in the sea ice community.

The addition of DCMU to the samples inhibited electron flow between PSII and PSI, thus preventing carbon fixation and the production of glucose. The subsequent drop in dissolved glucose concentrations thus potentially resulted from

both biotic and abiotic processes. However, in similar experiments with benthic microalgae, McMinn and Lee (2018) treated samples in the dark with antibiotics to inhibit bacterial activity. These samples showed no significant decline in glucose concentrations with time, indicating that abiotic breakdown of glucose was occurring at a much slower rate. Thus, bacterial consumption was responsible for most of the drop in glucose concentrations in the ice algae samples treated with DCMU. Glucose turnover times were between 2.9 and 17 h, rates consistent with open marine and freshwater turnover times elsewhere (Bunte and Simon 1999; Skoog et al. 2002; Alonso-Saez et al. 2012).

This study, together with that of McMinn and Lee (2018), has demonstrated that microsensors can be reliably used to measure microalgal exudation and bacterial uptake rates of glucose in these communities. Microsensors are well suited to these systems as biomass levels and DOC concentrations are often many times higher than in open water, which overcomes microsensor sensitivity issues.

Acknowledgements We would like to thank Antarctica NZ and the staff at Scott Base for providing support and logistics. We would also like to thank our reviewers, Nicole Hellesey and Katherina Petrou for their helpful and constructive criticisms.

Author contributions AMcM conceived the research, AM and FK conducted the field work, FK conducted the experiments, AMcM wrote the manuscript and all authors read and improved the manuscript.

Funding Open Access funding enabled and organized by CAUL and its Member Institutions. This research was supported by the New Zealand Antarctic Research Institute (NZARI Grant 2017-1-4) and the Australian Research Council's Special Research Initiative for Antarctic Gateway Partnership (Project ID SR140300001).

Data availability The authors declare that all data used in the production of this manuscript are presented in incorporated tables and also that all data support the claims made herein.

Declarations

Conflict of interest The authors are unaware of any conflicting interests. The authors have no relevant financial or non-financial interests to disclose.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Alonso-Sáez L, Sánchez O, Gasol JM (2012) Bacterial uptake of low molecular weight organics in the subtropical Atlantic: are major phylogenetic groups functionally different? *Limnol Oceanogr* 57:798–808
- Arrigo KR (2014) Sea ice ecosystems. *Ann Rev Mar Sci* 6:439–467
- Aslam SN, Cresswell-Maynard T, Thomas DN, Underwood GJC (2012) Production and characterization of the intra- and extracellular carbohydrates and polymeric substances (EPS) of three sea-ice diatom species, and evidence for a cryoprotective role for EPS. *J Phycol* 48:1494–1509
- Aslam SN, Michel C, Niemi A, Underwood GJC, Graham JC (2016) Patterns and drivers of carbohydrate budgets in ice algal assemblages from first year Arctic sea ice. *Limnol Oceanogr* 61:919–937
- Bellinger BJ, Underwood GJC, Ziegler SE, Gretz MR (2009) Significance of diatom-derived polymers in carbon flow dynamics within estuarine biofilms determined through isotopic enrichment. *Aquat Microb Ecol* 55:169–187
- Bunte C, Simon M (1999) Bacterioplankton turnover of dissolved free monosaccharides in a mesotrophic lake. *Limnol Oceanogr* 44:1862–1870
- Cook PLM, Revill AT, Butler ECV, Eyre BD (2004) Carbon and nitrogen cycling on intertidal mudflats of a temperate Australian estuary. II. Nitrogen cycling. *Mar Ecol Prog Ser* 280:39–54
- Cook PLM, Veuger B, Voer S, Middelburg JJ (2007) Effect of nutrient availability on carbon and nitrogen incorporation and flows through benthic algae and bacteria in near-shore sandy sediment. *Aquat Microb Ecol* 49:165–180
- Cummings VJ, Barr NG, Budd RD, Marriott PM, Safi KA (2019) Lohrer AM (2019) *In situ* response of Antarctic under-ice primary producers to experimentally altered pH. *Sci Rep* 9:6069. <https://doi.org/10.1038/s41598-019-42329-0>
- Fenchel T (2008) The microbial loop-25 years later. *J Exp Mar Biol Ecol* 366:99–103
- Genty B, Briantais J-M, Baker NR (1989) The relationship between quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochim Biophys Acta* 990:87–92
- Haas AF, Wild C (2010) Composition analysis of organic matter released by cosmopolitan coral reef-associated green algae. *Aquat Biol* 10:131–138
- Hama T, Yanagi K (2001) Production and neutral aldose composition of dissolved carbohydrates excreted by natural marine phytoplankton populations. *Limnol Oceanogr* 46:1945–1955
- Kennedy F, Martin A, McMinn A (2020) Insights into the production and role of nitric oxide in the Antarctic sea-ice diatom *Fragilariopsis cylindrus*. *J Phycol* 56:1196–1207
- Leles SG, Mitra A, Flynn KJ, Tilmann U, Stoecker D, Jeong HJ, Burkholder J, Hansen PJ, Caron DA, Gilbert PM, Hallegraeff GM, Raven JA, Sanders RW, Zubkov M (2019) Sampling bias misrepresents the biogeographical significance of constitutive mixotrophs across global oceans. *Global Ecol Biogeogr* 28:418–428
- Lizotte MP (2001) The contributions of sea ice algae to Antarctic marine primary production. *Amer Zool* 41:57–73
- Martin A, Anderson MJ, Thorn C, Davy SK, Ryan KG (2011) Response of sea ice microbial communities to environmental change: an in situ experiment in the Antarctic. *Mar Ecol Prog Ser* 424:25–37
- Martin A, McMinn A, Davy SK, Miller HC, Anderson MJ, Hall JA, Ryan KG (2012) Preliminary evidence for the microbial loop in Antarctic sea ice using microcosm simulations. *Antarct Sci* 24:547–553
- McMinn A, Lee S (2018) Use of glucose biosensors to measure extracellular glucose production by intertidal microphytobenthos in southern Tasmania. *J Phycol* 54:410–418
- McMinn A, Gibson J, Hodgson D, Aschman J (1995) Nutrient limitation in Ellis Fjord, Antarctica. *Pol Biol* 15:269–276
- McMinn A, Skerratt J, Trull T, Ashworth C, Lizotte M (1999) Nutrient stress gradient in the bottom 5 cm of fast ice, McMurdo sound, Antarctica. *Pol Biol* 21:220–227
- McMinn A, Ryan K, Gademann R (2003) Photoacclimation of Antarctic fast ice algal communities determined by pulse amplitude modulation (PAM) fluorometry. *Marine Biology* 143:359–367
- McMinn A, Ryan KR, Ralph PJ, Pankowski A (2007) Spring sea ice photosynthesis, primary productivity and biomass distribution in eastern Antarctica, 2002–2004. *Marine Biology* 151:985–995
- McMinn A, Pankowski A, Ashworth C, Bhagooli R, Ralph P, Ryan K (2010) In situ net primary productivity and photosynthesis of Antarctic sea ice algal, phytoplankton and benthic algal communities. *Marine Biology* 157:1345–1356. <https://doi.org/10.1007/s00227-010-1414-8>
- McMinn A, Ashworth C, Bhagooli R, Martin A, Salleh S, Ralph P, Ryan K (2012) Antarctic coastal microalgal primary production and photosynthesis. *Mar Biol* 159:2827–2837
- McMinn A, Liang Y, Wang M (2020) Minireview: the role of viruses in marine photosynthetic biofilms. *J Mar Life Sci Technol* 2:203–208
- Myklesstad S (1977) Production of carbohydrates by marine planktonic diatoms. II influence of the N/P ratio in the growth medium on the assimilation ratio, growth rate and production of extracellular carbohydrates by *Chaetoceros affinis* var *williei* (Gran) Hustedt and *Skeletonema costatum* (Grev) Cleve. *J Exp Mar Biol Ecol* 9:125–136
- Oakes JM, Eyre BD, Middelburg JJ, Boschker HTS (2010) Composition, production, and loss of carbohydrates in subtropical shallow subtidal sandy sediments: rapid processing and long-term retention revealed by C-13-labeling. *Limnol Oceanogr* 55:2126–2138
- Palmisano AC, Sullivan CW (1985) Pathways of photosynthetic carbon assimilation in sea ice microalgae from McMurdo Sound, Antarctica. *Limnol Oceanogr* 30:674–678
- Rivaro P, Ardini F, Grotti M, Vivado D, Salis A, Damonte G (2021) Detection of carbohydrates in sea ice extracellular polymeric substances via solid-phase extraction and HPLC-ESI-MS/MS. *Mar Chem* 228:103911
- RStudio Team (2021) RStudio: integrated development environment for R. RStudio, Boston
- Skoog A, Whitehead K, Sperling F, Junge K (2002) Microbial glucose uptake and growth along a horizontal nutrient gradient in the North Pacific. *Limnol Oceanogr* 47:1676–1683
- Smith DJ, Underwood GJC (2000) The production of extracellular carbohydrate exopolymers (EPS) by estuarine benthic diatoms: the effects of growth phase and light and dark treatment. *J Phycol* 36:321–333
- Sorrell BK, Hawes I, Stratmann T, Lund-Hansen LC (2021) Photobiological effects on ice algae of a rapid whole-fjord loss of snow cover during spring growth in Kangerlussuaq, a West Greenland Fjord. *J Mar Sci Eng* 9:814
- Staats N, Stal J, Mur R (1999) Oxygenic photosynthesis as driving process in exopolysaccharide production of benthic diatoms. *Mar Ecol Prog Ser* 193:261–269
- Staats N, Stal LJ, Mur LR (2000) Oxygenic photosynthesis as driving process in exopolysaccharide production of benthic diatoms. *Mar Ecol Prog Ser* 193:261–269
- Ugalde SC, Martin A, Meiners KM, McMinn A, Ryan KG (2014) Extracellular organic carbon dynamics during a bottom-ice algal bloom (Antarctica). *Aquat Microb Ecol* 73:195–210
- Underwood GJC, Kromkamp J (1999) Primary production by phytoplankton and microphytobenthos in estuaries. *Adv Ecol Res* 29:93–153
- Underwood GJC, Paterson DM (2003) The importance of extracellular carbohydrate production by marine epipelagic diatoms. *Adv Bot Res* 40:183–240
- Underwood GJC, Fietz S, Papadimitriou S, Thomas DN, Dieckmann GS (2010) Distribution and composition of dissolved extracellular polymeric substances (EPS) in Antarctic sea ice. *Mar Ecol Prog Ser* 404:1–19

- Underwood GJC, Aslam SN, Michel C, Niemi A, Norman L, Meiners KM, Laybourn-Parry J, Paterson H, Thomas DN (2013) Broad-scale predictability of carbohydrates and exopolymers in Antarctic and Arctic sea ice. *PNAS* 110:15734–15739
- Van Leeuwe MA, Tedesco L, Arrigo KR, Assmy P, Campbell K, Meiners KM, Rintalla J-M, Selz V, Thomas DN, Stefels J (2018) Microalgal community structure and primary production in Arctic and Antarctic sea ice: a synthesis. *Elementa* 6:4. <https://doi.org/10.1525/elementa.267>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.